

Extrapankreatic actions of incretin-based therapies on bone in diabetes mellitus

Sity Aishah Mansur, BSc (Hons), MEng

Faculty of Life and Health Sciences of Ulster University

Thesis submitted for the degree of Doctor of Philosophy

October 2015

I confirm that the word count of this thesis is less than 100,000 words excluding the title page, contents acknowledgements, summary or abstract, abbreviations, footnotes, diagrams, maps, illustrations, tables, appendices, and references or bibliography

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	xiii
SUMMARY	xiv
ABBREVIATIONS	xvi
DECLARATION	xix

CHAPTER 1

GENERAL INTRODUCTION

1.0 OVERVIEW	2
1.1 DIABETES MELLITUS	2
1.1.1 Type 1 diabetes mellitus (T1DM)	3
1.1.2 Type 2 diabetes mellitus (T2DM)	4
1.1.3 Existing treatment for diabetes	5
1.2 BONE	10
1.2.1 Assessment of bone mass	12
1.2.2 Bone microarchitecture	13
1.2.3 Biomechanical test	14
1.2.4 Bone material properties	14
1.3 EFFECTS OF DIABETES ON THE SKELETON	15
1.3.1 Hyperglycaemia and AGEs	15
1.3.2 Insulin	16
1.3.3 Growth factors and cytokines	17
1.3.4 Bone impairments in animal models of diabetes	18
1.3.5 Bone mineral density and fracture risk in human diabetes	20

1.4	INCRETIN EFFECT	21
1.5	GLUCOSE-DEPENDENT INSULINOTROPIC POLYPEPTIDE (GIP)	22
1.5.1	Synthesis, secretion and signaling	22
1.5.2	Pancreatic actions	23
1.5.3	Extrapancreatic actions	23
1.5.4	DPP-4 and GIP metabolism	25
1.5.5	Structurally-modified stable GIP agonists	26
1.6	GLUCAGON-LIKE PEPTIDE-1 (GLP-1)	27
1.6.1	Synthesis, secretion and signaling	27
1.6.2	Pancreatic actions	29
1.6.3	Extrapancreatic actions	30
1.6.4	DPP-4 and GLP-1 metabolism	31
1.6.5	Structurally-modified stable GLP-1 agonists	32
1.6.6	Exendin-4	32
1.6.7	Liraglutide	33
1.6.8	Newly-approved/developed GLP-1 therapeutics	33
1.7	DPP-4 INHIBITORS	35
1.8	FUNCTION OF INCRETINS ON BONE	36
1.9	AIMS OF THESIS	39

CHAPTER 2

GENERAL MATERIALS AND METHODS

2.1	PEPTIDES	41
2.1.1	Synthesis of peptides	41
2.1.2	Identification and characterization	41

2.2	CELL CULTURE	41
2.2.1	SaOS-2 cells	41
2.3	ANIMALS	42
2.3.1	GIPR KO mice	42
2.3.2	GLP-1-R KO mice	42
2.3.3	Double Incretin receptor (DIR) KO mice	43
2.3.4	NIH Swiss <i>TO</i> mice	43
2.3.5	C57 BL/KsJ diabetic (<i>db/db</i>) mice	43
2.3.6	High-fat fed NIH Swiss mice	43
2.3.7	Collection of blood samples	44
2.3.8	Intraperitoneal glucose tolerance test	44
2.3.9	Intraperitoneal insulin sensitivity test	44
2.4	<i>IN VIVO</i> BIOCHEMICAL ANALYSES	44
2.4.1	Plasma glucose determination	44
2.4.2	Plasma insulin determination	45
2.5	MEASUREMENT OF BODY COMPOSITION, BONE MINERAL DENSITY AND MINERAL CONTENT BY DUAL-ENERGY X-RAY ABSORPTIOMETRY (DEXA) SCANNING	47
2.6	ASSESSMENT ON BONE STRENGTH AND QUALITY	48
2.6.1	Quantitative x-ray imaging (qXRI)	49
2.6.2	X-ray microcomputed tomography (microCT)	51
2.6.3	Three-point bending test	54
2.6.4	Nanoindentation	57
2.6.5	Quantitative backscattered electron imaging (qBEI)	59
2.7	STATISTICAL ANALYSIS	62

CHAPTER 3

EFFECT OF GIP PEPTIDES ON SAOS-2 CELLS *IN VITRO* AND GENETIC DEFICIENCIES OF GIP AND/OR GLP-1 RECEPTORS ON BONE MASS *IN VIVO*

3.1	SUMMARY	64
3.2	INTRODUCTION	65
3.3	MATERIALS AND METHODS	67
3.3.1	Peptides	67
3.3.2	Maintenance of SaOS-2 cells	67
3.3.3	Animals and study design	67
3.3.3.1	GIPR KO mice	67
3.3.3.2	GLP-1R KO mice	67
3.3.3.3	Double Incretin receptor (DIR) KO mice	68
3.3.4	Measurement of TGF- β	68
3.3.5	Measurement of IGF-1	68
3.3.6	Measurement of bone alkaline phosphatase activity	
3.3.6.1	Alkaline phosphatase determination	69
3.3.6.2	Total protein determination	70
3.3.7	Measurement of cyclic AMP	70
3.3.8	Measurement of plasma glucose concentration	70
3.3.9	Measurement of body composition, bone density and mineral content by DEXA scanning	71
3.3.10	Statistical analysis	71
3.4	RESULTS	71
3.4.1	Dose-dependent effects of native GIP and [D- Ala ²]GIP on TGF- β 1 and IGF-1 release from SaOS- 2 cells	71

3.4.2	Dose-dependent effects of native GIP and [D-Ala ²]GIP on alkaline phosphatase activity in SaOS-2 cells	72
3.4.3	Dose-dependent effects of native GIP and [D-Ala ²]GIP on cAMP production in SaOS-2 cells	72
3.4.4	Comparison of body composition in GIPR, GLP-1R, DIR KO and normal mice	72
3.4.5	Comparison of glucose tolerance in GIPR, GLP-1R, DIR KO and normal mice	73
3.4.6	Comparison of bone mineral density and mineral content in GIPR, GLP-1R, DIR KO and normal mice	73
3.5	DISCUSSION	74

CHAPTER 4

RECOVERY OF CORTICAL BONE STRENGTH AT THE TISSUE LEVEL BY [D-ALA²]GIP AND LIRAGLUTIDE IN STREPTOZOTOCIN-INDUCED INSULIN-DEFICIENT DIABETIC MICE

4.1	SUMMARY	95
4.2	INTRODUCTION	96
4.3	MATERIALS AND METHODS	98
4.3.1	Synthesis of [D-Ala ²]GIP and Liraglutide	98
4.3.2	Animals and study design	98
4.3.3	Measurement of plasma glucose and insulin	99
4.3.4	Measurement of body composition, bone density and mineral content by DEXA scanning	99
4.3.5	Assessment of bone quality and strength	99

4.3.5.1	Quantitative X-ray Imaging (qXRI)	99
4.3.5.2	X-ray microcomputed tomography (μ CT)	99
4.3.5.3	Three-point bending test	100
4.3.5.4	Nanoindentation	100
4.3.5.5	Quantitative backscattered electron imaging (qBEI)	100
4.3.6	Statistical analysis	100
4.4	RESULTS	101
4.4.1	Effects of once-daily administration of [D-Ala ²]GIP or Liraglutide on metabolic parameters in streptozotocin-induced diabetic mice	101
4.4.2	Effects of once-daily administration of [D-Ala ²]GIP or Liraglutide on glucose tolerance and glucose-induced plasma insulin release in streptozotocin-induced diabetic mice	101
4.4.3	Effects of once-daily administration of [D-Ala ²]GIP or Liraglutide on insulin sensitivity in streptozotocin-induced diabetic mice	102
4.4.4	Effects of [D-Ala ²]GIP and Liraglutide on total body fat, bone mineral density (BMD) and bone mineral content (BMC) in streptozotocin-induced diabetic mice	102
4.4.5	Effects of [D-Ala ²]GIP and Liraglutide on femoral mineral content in streptozotocin-induced diabetic mice	102
4.4.6	Effects of [D-Ala ²]GIP and Liraglutide on trabecular bone morphology and cortical bone geometry in streptozotocin-induced diabetic mice	103
4.4.7	Effects of [D-Ala ²]GIP and Liraglutide on whole-bone mechanical properties in streptozotocin-induced diabetic mice	103

4.4.8	Effects of [D-Ala ²]GIP and Liraglutide on nanomechanical properties of cortical bone matrix in streptozotocin-induced diabetic mice	104
4.4.9	Effects of [D-Ala ²]GIP and Liraglutide on bone mineral density distribution in streptozotocin-induced diabetic mice	104
4.5	DISCUSSION	105

CHAPTER 5

SIMULTANEOUS ACTIVATION OF GIP, GLP-1 AND GLUCAGON RECEPTORS BY [D-ALA²]GIP-OXYNTOMODULIN POSITIVELY AFFECTS CORTICAL AND TRABECULAR BONE PROPERTIES AND IMPROVES BONE MECHANICAL STRENGTH IN GENETICALLY-INDUCED C57 BL/KsJ DIABETIC (DB/DB) MICE

5.1	SUMMARY	138
5.2	INTRODUCTION	139
5.3	MATERIALS AND METHODS	141
5.3.1	Synthesis of [D-Ala ²]GIP-Oxyntomodulin	141
5.3.2	Animals and study design	141
5.3.3	Measurement of plasma glucose and insulin	141
5.3.4	Measurement of body composition, bone density and mineral content by DEXA scanning	141
5.3.5	Assessment of bone quality and strength	142
5.3.5.1	Quantitative X-ray Imaging (qXRI)	142
5.3.5.2	X-ray microcomputed tomography (μCT)	
5.3.5.3	Three-point bending test	142

5.3.5.4 Nanoindentation	143
5.3.5.5 Quantitative backscattered electron imaging (qBEI)	143
5.3.6 Statistical analysis	143
5.4 RESULTS	144
5.4.1 Effects of once-daily administration of [D-Ala ²]GIP-Oxm on metabolic parameters in <i>db/db</i> mice	144
5.4.2 Effects of once-daily administration of [D-Ala ²]GIP-Oxm on glucose tolerance, glucose-induced plasma insulin release and insulin sensitivity in <i>db/db</i> diabetic mice	144
5.4.3 Effects of [D-Ala ²]GIP-Oxm on percentage body fat, bone mineral density and bone mineral content in <i>db/db</i> mice	144
5.4.4 Effects of [D-Ala ²]GIP-Oxm on cortical and trabecular bone mineral content in <i>db/db</i> mice	145
5.4.5 Effects of [D-Ala ²]GIP-Oxm on trabecular bone microarchitecture and cortical bone geometry in <i>db/db</i> mice	145
5.4.6 Effects of [D-Ala ²]GIP-Oxm on whole-bone strength in <i>db/db</i> mice	146
5.4.7 Effects of [D-Ala ²]GIP-Oxm on cortical bone nanomechanical properties in <i>db/db</i> mice	146
5.4.8 Effects of [D-Ala ²]GIP-Oxm on cortical bone mineral density distribution in <i>db/db</i> mice	147
5.5 DISCUSSION	148

CHAPTER 6

EFFECTS OF THE GLP-1R AGONIST EXENDIN-4 AND THE DPP-4-INHIBITOR SITAGLIPTIN ON BONE STRENGTH AND QUALITY IN HIGH-FAT FED MICE

6.1	SUMMARY	179
6.2	INTRODUCTION	180
6.3	MATERIALS AND METHODS	182
6.3.1	Animals and study design	182
6.3.2	Measurement of body composition, bone density and mineral content by DEXA scanning	182
6.3.3	Assessment of bone mineral content by quantitative x-ray imaging (qXRI)	183
6.3.4	Assessment of trabecular bone morphology and cortical geometry by X-ray microcomputed tomography (microCT)	183
6.3.5	Evaluation of extrinsic mechanical strength of bones	183
6.3.6	Statistical analysis	183
6.4	RESULTS	184
6.4.1	Effects of twice-daily administration of Exendin-4 on metabolic parameters, glucose tolerance and insulin sensitivity in high-fat fed NIH Swiss mice	184
6.4.2	Effects of once-daily administration of Sitagliptin on metabolic parameters, glucose tolerance and insulin sensitivity in high-fat fed mice NIH Swiss mice	184
6.4.3	Effects of Exendin-4 and Sitagliptin on total body fat mass, bone mineral density and bone mineral content in high-fat fed NIH Swiss mice	184

6.4.4	Effects of Exendin-4 and Sitagliptin on cortical and trabecular bone mineral content in high-fat fed NIH Swiss mice	185
6.4.5	Effects of Exendin-4 and Sitagliptin on trabecular microstructural morphology and cortical bone geometry in high-fat fed NIH Swiss mice	185
6.4.6	Effects of Exendin-4 and Sitagliptin on mechanical properties of cortical bone in high-fat fed NIH Swiss mice	186
6.5	DISCUSSION	187

CHAPTER 7

GENERAL DISCUSSION

7.1	INTRODUCTION	225
7.2	<i>IN VITRO</i> STUDIES WITH GIP	226
7.3	ACTIVATION OF GIPR OR GLP-1R ON BONE QUALITY IN PHARMACOLOGICALLY-INDUCED INSULIN-DEFICIENT DIABETIC MICE	227
7.4	SIMULTANEOUS ACTIVATION OF GIPR AND GLP-1R BY [D-ALA²]GIP-OXM ON BONE QUALITY IN LEPTIN RECEPTOR-DEFICIENT C57 BL/KsJ (<i>DB/DB</i>) MICE	229
7.5	EFFECTS OF CURRENT INCRETIN-BASED THERAPIES FOR HUMAN T2DM ON BONE QUALITY IN HIGH-FAT FED MICE	230

7.6	STRENGTHS AND WEAKNESSES	231
------------	---------------------------------	------------

7.7	SUMMARY OF FUTURE STUDIES	233
------------	----------------------------------	------------

CHAPTER 8

REFERENCES	235
-------------------	------------



ACKNOWLEDGEMENTS

First and foremost, I would like to express my deepest gratitude to Allah for granting me good health (Alhamdulillah) and allowing me to successfully finish my research. Thank you to Malaysian government and Universiti Tun Hussein Onn Malaysia (UTHM) for giving me the opportunity to further my postgraduate study abroad and supporting me financially.

I would also like to offer my thanks to my 1st supervisor, Dr. Nigel Irwin for the hard work and excellent advice throughout the three years of my PhD. For my 2nd supervisor, Professor Peter Flatt, thanks for your guidance and support. Massive thank you for Dr. Guillaume Mabilieu from University of Angers, France for his active participation in my research and helping me with bone quality assessment for the long-term animal studies in the thesis.

This dissertation is dedicated to my family, especially my husband, Adri and my son, Aydeen. Without their unconditional love and constant encouragement, this thesis would not be possible. For those in Malaysia : mak, babah, mama, my sisters, brothers and not forgotten, my three bestfriends; Dido, Aju and June, thanks for your continuous prayers and for believing in me. Finally, I would like to thank all the members of Diabetes Research Group for their help and friendship, which makes this 3-year journey a meaningful one.

SUMMARY

Diabetes mellitus is correlated with modifications in bone microarchitectural and mechanical strength, leading to increased bone fragility. The incretin hormones, with a classical effect to increase insulin secretion following food ingestion, are now postulated to have important direct effects on bone. As such, glucose-dependent insulintropic polypeptide (GIP) has dual actions on bone cells; enhancing bone-forming activity of osteoblasts and suppressing bone resorption by osteoclasts. The sister incretin of GIP, glucagon-like peptide-1 (GLP-1), is also suspected to directly influence bone health in a beneficial manner, although mechanism are less clear at present. The physiological actions of incretins are attenuated by dipeptidyl peptidase (DPP-4) activity and it is speculated that introduction of DPP-4 inhibitor may also positively affect quality of the skeleton. As such, this thesis evaluates the potential beneficial effects of a DPP-4 resistant GIP analogue, namely [D-Ala²]GIP, on osteoblastic-derived, SaOS-2 cells, and also preliminary *in vivo* studies on the impact of genetic deficiencies of GIPRs and GLP-1Rs on bone mineral density and content. Further studies characterised the beneficial effects of incretin-based therapies on metabolic control, bone microstructure and bone mechanical integrity in animal models of pharmacologically-, genetically- and environmentally-induced diabetes. GIP and related stable analogue increased bone-forming biomarkers in SaOS-2 cells and importantly, [D-Ala²]GIP was shown to be more potent than native GIP. Knockout mouse studies revealed that both GIPR and GLP-1R signaling are important for optimum bone mass. All diabetic mouse models displayed reduced bone mass, altered bone micromorphology and impairment of bone mechanical strength, similar to the human situation, confirming their appropriateness. The incretin-based therapeutics, [D-Ala²]GIP and Liraglutide, in streptozotocin-diabetic significantly increased bone matrix properties, indicating recovery of bone strength at the tissue level. The beneficial effects of administration of [D-Ala²]GIP-oxynatomodulin on bone health in *db/db* mice were more prominent as the Oxm analogue did not only improve bone strength at tissue level, but also at whole-bone level. These modifications were independent of metabolic status. Twice-daily Exendin-4 therapy improved glycaemic control and increased work required to resist bone fracture in high-fat fed mice. It was also established that Sitagliptin had neutral effects on bone microstructure and mechanical strength in high-fat mice. In

summary, these data demonstrate the negative impact of diabetes mellitus on normal skeleton development and bone quality. Moreover, this thesis highlights the growing potential of incretin-based therapies for ameliorating bone defects and improving the increased fragility fracture risk associated with diabetes.



ABBREVIATIONS

AGEs	Advanced glycation end products
AlkP	Alkaline phosphatase
AMD	Absorbing mineral density
AUC	Area under the curve
B.Dm	Bone diameter
BMC	Bone mineral content
BMD	Bone mineral density
BMDD	Bone mineral density distribution
BSE	Back-scattered electron
BV	Bone volume
bw	Body weight
cAMP	Cyclic adenosine monophosphate
cm	Centimeter
CSMI	Cross-sectional moment of inertia
Ct.Th	Cortical thickness
(db/db)	Diabetic (db/db) mice
DEXA	Dual-energy x-ray absorptiometry
dl	Decilitre
DM	Diabetes mellitus
FBS	Fetal bovine serum
g	Gram
GIP	Glucose-dependent insulintropic polypeptide
GIPR	Glucose-dependent insulintropic polypeptide receptor
GL	Grey level
GLP-1	Glucagon-like peptide-1
GLP-1R	Glucagon-like peptide-1 receptor
GPa	Gigapascal
h	Hour
ip	Intraperitoneal injection
keV	Kilo electron volt
kg	Kilogram
KO	Knock-out mice

kV	Kilovolt
l	Litre
Ma.Dm	Marrow diameter
MALDI-TOF-MS	Matrix-assisted laser desorption/ionization-time of flight mass spectrometry
mg	Miligram
microCT	X-ray microcomputed tomography
min	Minutes
ml	Mililitre
mm	Milimeter
mmol	Milimolar
MMP	Matrix metalloproteinase
mN	Milnewton
MPa	Megapascal
MSC	Mesenchymal stem cell
N	Newton
nm	Nanometer
OPG	Osteoprotegerin
Oxm	Oxyntomodulin
pA	Picoampere
pJ	Picojoule
PMMA	Polymethylmethacrylate
qBEI	Quantitative back-scattered electron imaging
qXRI	Quantitative x-ray imaging
RANKL	Receptor activator on nuclear factor-Kappa B ligand
RIA	Radioimmunoassay
ROI	Region of interest
rpm	Revolutions per minute
RT-PCR	Reverse transcriptase-polymerase chain reaction
SEM	Standard error of the mean
STZ	Streptozotocin
Tb.N	Trabecular number
Tb.Sp	Trabecular separation
Tb.Th	Trabecular thickness

TRAP	Tartrate-resistant acid phosphatase
TV	Trabecular volume
U	Unit
Z	Atomic number
μA	Microampere
μg	Microgram
μl	Microlitre
μm	Micrometer



DECLARATION

“I hereby declare that for 2 years following the date on which the thesis is deposited in Research Student Administration of Ulster University, the thesis shall remain confidential with access or copying prohibited. Following expiry of this period I permit

1. the Librarian of the University to allow the thesis to be copied in whole or in part without reference to me on the understanding that such authority applies to the provision of single copies made for study purposes or for inclusion within the stock of another library.
2. the thesis to be made available through the Ulster Institutional Repository and/or EThOS under the terms of the Ulster eTheses Deposit Agreement which I have signed.

IT IS A CONDITION OF USE OF THIS THESIS THAT ANYONE WHO CONSULTS IT MUST RECOGNISE THAT THE COPYRIGHT RESTS WITH THE UNIVERSITY AND THEN SUBSEQUENTLY TO THE AUTHOR ON THE EXPIRY OF THIS PERIOD AND THAT NO QUOTATION FROM THE THESIS AND NO INFORMATION DERIVED FROM IT MAY BE PUBLISHED UNLESS THE SOURCE IS PROPERLY ACKNOWLEDGED.”

Chapter 1

General Introduction



PTTHM
PERPUSTAKAAN TUNKU TUN AMINAH

1.0 OVERVIEW

Diabetes has become a global issue as it is estimated that more than 387 million people worldwide are affected by this modern disease (International Diabetes Federation 2014b). The alarming number of diabetes cases has listed this disorder as the potential seventh leading cause of death in 2030 (Mathers & Loncar 2006). Common complications that are associated with diabetes include retinopathy (blindness), neuropathy (nerve damage), cardiovascular disease, stroke and nephropathy (kidney failure) (Diabetes UK 2014) and most recently, diabetes is also related to bone loss. Extensive studies on the impact of diabetes on bone have linked the disease to severe impairment of bone mechanical integrity, reduced bone mineral density (BMD) in T1DM and unaffected or even increased BMD in T2DM. Regardless of the inconsistencies in BMD, both types of diabetic patients have increased risk of fractures (Janghorbani et al. 2007, Vestergaard 2007). Early diagnosis and treatment of diabetes is vital to prevent long term complications that could lead to bone fractures and mortality in elderly people. Furthermore, the link between diabetes and bone fragility suggests that if blood glucose management is improved or if signaling pathways contributing to diabetic bone pathologies are corrected, overall bone quality could be enhanced, improving the increased risk of fracture in diabetic patients.

1.1 DIABETES MELLITUS

Diabetes mellitus (DM) is a metabolic disease where production of insulin is insufficient or body cannot effectively utilise insulin produced by pancreas. As a consequence, there will be high blood glucose levels in the body. Diagnosis of diabetes mellitus is made when fasting plasma glucose is ≥ 126 mg/dl (7.0 mmol/l) or plasma glucose level is ≥ 200 mg/dl (11.1 mmol/l) measured 2 hours after 75 g-oral glucose tolerance test (Zemmit et al. 2001). Generally, the disease is classified into 2 major forms; type 1 diabetes mellitus (T1DM) and type 2 diabetes mellitus (T2DM).

1.1.1 Type 1 diabetes mellitus

Type 1 diabetes mellitus (T1DM), previously referred to as juvenile-onset or insulin-dependent diabetes, is characterised by inability of the body to synthesise insulin due to destruction of insulin-producing pancreatic beta cells. This results in a dependency on exogenous insulin to stabilise blood glucose concentrations following feeding. The pathogenesis of this autoimmune disorder has been extensively studied in rodent models and humans and several key target for autoantibodies and autoreactive cells have been identified; insulin, glutamic acid decarboxylase (GAD), protein tyrosine phosphatase insulinoma antigen (IA-2), zinc transporter 8 (ZnT8) and islet glucose-6-phosphatase catalytic subunit-related protein (IGRP) (Culina et al. 2013, Yoon & Jun 2005, Zhang et al. 2008). Pancreatic beta cells autoantigens together with lymphocytes and macrophages trigger overexpression of CD4⁺ and beta cell-specific cytotoxic CD8⁺ T cells (Pinkse et al. 2005) which eventually lead to the destruction of insulin-producing beta cells.

The primary cause of T1DM is autoimmune beta-cell destruction. However, there is also correlation between genetic and several environmental factors in the pathogenesis of T1DM. Genetic components that strongly influence T1DM are reported to be on major histocompatibility complex human leukocyte antigen (HLA) alleles, which are associated with 'protection from' or 'susceptibility towards' T1DM. More than 90% of T1DM patients have either DR3-DQ2 or DR4-DQ8 haplotypes, while the percentage of normal controls with these haplotypes is less than 40% (Atkinson & Eisenbarth 2001). These genes are known to be important regulators in immune response by introducing peptide antigens to T lymphocytes which stimulate the production of antibodies against beta-cell antigens. Further exposure to environmental triggers such as viruses in susceptible individuals can cause abnormal activation of immune system that will generate multiple autoantibodies to islet cells.

T1DM is most prevalent in children and young people and it accounts for about 10% of diabetes cases worldwide (International Diabetes Federation 2014a). Statistical studies conducted by EURODIAB (EUROpe and DIABetes) from 1989 to 2003 estimated that diabetes cases in children below 15 years old will rise by 70% in 2020

(Patterson et al. 2009). A more recent study on trends over 20-year incidence projected that the prevalence of Type 1 diabetes has annual increase of 3.4% (Patterson et al. 2012).

1.1.2 Type 2 diabetes mellitus

Type 2 diabetes mellitus (T2DM), previously known as adult-onset or non-insulin-dependent diabetes, develops when beta cells do not make enough insulin or the insulin produced does not work appropriately. T2DM is a progressive disease and starts gradually with insulin insensitivity or impairment of the body's response to insulin (Stumvoll et al. 2005). Compensatory mechanism in the body respond to insulin resistance by secreting more insulin to maintain normal glucose levels, but eventually pancreatic beta-cells can no longer produce sufficient insulin to maintain normal glucose levels. Over time, worsening insulin resistance and the decline in beta cell function will lead to hyperglycaemia and eventually progress to the diabetic state. The environmental factors that trigger T2DM are pregnancy, ageing and modern lifestyle (high calorie intake, obesity, physical inactivity) (Chen et al. 1988, Hu 2011, Stumvoll et al. 2005). Similar to T1DM, the combination of genetic predisposition and environmental factors contributes to the onset of insulin resistance and its progression to T2DM (Leahy 2005). The key genes associated with genetic predisposition of T2DM are *TCF7L2*, *PPARG*, *KCNJ11*, *CAPN10*, *HHEX/IDE*, *MC4R*, *FTO* and *KCNQ1* (Tahrani et al. 2011, Valeriya et al. 2007).

T2DM usually appears in middle-aged and older people and accounts for 90% of all diabetes cases (International Diabetes Federation 2014a). T2DM has become an epidemic as this disease progresses slowly and undetected many years before diagnosis is made. With increasing prevalence of obesity and T2DM, a huge burden is imposed on the public health-system (Diabetes UK 2012).

1.1.3 Existing treatment for diabetes

The only therapeutic option for T1DM patients is exogenous insulin replacement therapy. However, T1DM complications are not always prevented by exogenous insulin therapy (Atkinson et al. 2014). Other important glucoregulatory hormone which is co-secreted with insulin by pancreatic beta cells, amylin, is also reduced in T1DM, leading to generation of synthetic amylin analogue, Pramlintide. Amylin replacement with Pramlintide improves glycaemic control by delaying gastric emptying and suppressing postprandial glucagon secretion (Fineman et al. 2002, Nyholm et al. 1999). In a long term clinical study, adjunct therapy with Pramlintide has been shown to provide better glycaemic control in T1DM individuals compared to insulin replacement alone (Ratner et al. 2004). As such, this amylin-related analogue was clinically approved in April 2004 by the US Food and Drug Administration (FDA) to be used as add-on with insulin therapy in T1DM patients (Ryan et al. 2005). Another option for T1DM treatment is islet-cell transplantation, leaving the recipients dependent on lifelong immunosuppression drugs to prevent organ rejection. Most importantly, the limiting factor in this type of treatment is a short supply of organs and until new sources of functional islets are found, the future for organ transplantation in T1DM treatment remains sceptical (McCall & Shapiro 2012).

Obesity is normally associated with impaired glucose tolerance and increased risk of developing T2DM. This is due to increased adipocyte accumulation of non-esterified fatty acids, hormones, inflammatory cytokines and other molecules that are involved in the development of insulin resistance (Kahn et al. 2006). Essentially, insulin resistance is improved through intensive lifestyle intervention by monitoring nutrient intake and regular physical activity (Tahrani et al. 2011). However, lifestyle changes are difficult to maintain and not sufficient to keep blood glucose levels normal. Consequently, oral or injectable antidiabetic drugs are often required to manage T2DM (International Diabetes Federation 2014a). Currently available T2DM treatments are summarised in Table 1.1.

Chapter 8

References



PTTAUTHM
PERPUSTAKAAN TUNKU TUN AMINAH

Agerso H, Jensen LB, Elbrond B, Rolan P & Zdravkovic M 2002 The pharmacokinetics, pharmacodynamics, safety and tolerability of NN2211, a new long-acting GLP-1 derivative, in healthy men. *Diabetologia* **45** 195-202.

Ahren B 2007 GLP-1-based therapy of type 2 diabetes : GLP-1 mimetics and DPP-IV inhibitors. *Current Diabetes Reports* **7** 340-347.

Akune T, Ohba S, Kamekura S, Yamaguchi M, Chung U, Kubota N, Terauchi Y, Harada Y, Azuma Y, Nakamura K, Kadowaki T & Kawaguchi H 2004 PPAR- γ insufficiency enhances osteogenesis through osteoblast formation from bone marrow progenitors. *Journal of Clinical Investigation* **113** 846-855.

Alexopoulou O, Jamart J, Devogelaer JP, Brichard S, de Nayer P & Buysschaert M 2006 Bone density and markers of bone remodeling in type 1 male diabetic patients. *Diabetes & Metabolism* **32** 453-458.

Alikhani M, Alikhani Z, Boyd C, MacLellan CM, Raptis M, Liu R, Pischon N, Trackman PC, Gerstenfeld L & Graves DT 2007 Advanced glycation end products stimulate osteoblast apoptosis via the MAP kinase and cytosolic apoptotic pathways. *Bone* **40** 345-353.

Aschner P, Kipnes MS, Luncford JK, Sanchez M, Mickel C & Williams-Herman DE 2006 Effect of the dipeptidyl peptidase-4 inhibitor Sitagliptin as monotherapy on glycemic control in patients with type 2 diabetes. *Diabetes care* **29** 2632-2637.

Ashcroft FM 1996 Mechanisms of the glycaemic effects of sulfonylureas. *Hormone and Metabolic Research* **28** 456-463.

Atkinson MA & Eisenbarth GS 2001 Type 1 diabetes: new perspectives on disease pathogenesis and treatment. *The Lancet* **358** 221-229.

Atkinson MA, Eisenbarth GS & Michels AW 2014 Type 1 diabetes. *Lancet* **383** 69-82.

- Baggio LL, Huang Q, Brown TJ & Drucker DJ 2004 Oxyntomodulin and glucagon-like peptide-1 differentially regulate murine food intake and energy expenditure. *Gastroenterology* **127** 546-558.
- Baggio LL & Drucker DJ 2007 Biology of Incretins: GLP-1 and GIP. *Gastroenterology* **132** 2131-2157.
- Barnett A 2006 DPP-4 inhibitors and their potential role in the management of type 2 diabetes. *International Journal of Clinical Practice* **60** 1454-1470.
- Barnett AH 2011 Lixisenatide: evidence for its potential use in the treatment of type 2 diabetes. *Core Evidence* **6** 67-79.
- Bassett JHD, Van der Spek A, Gogakos A & Williams GR 2012 Quantitative x-ray imaging of rodent bone by faxitron. *Bone Research Protocols, Methods in Molecular Biology* **816** 499-506.
- Bell GI, Santerre RF & Mullenbach GT 1983 Hamster preproglucagon contains the sequence of glucagon and two related peptides. *Nature* **302** 716-718.
- Bergental RM, Wysham C, MacConell L, Malloy J, Walsh B, Yan P, Wilhelm K, Malone J & Porter LE 2010 Efficacy and safety of exenatide once weekly versus sitagliptin or pioglitazone as an adjunct to metformin for treatment of type 2 diabetes (DURATION-2): a randomised trial. *Lancet* **376** 431-39.
- Bhat VK, Kerr BD, Flatt PR & Gault VA 2013 A novel GIP-oxyntomodulin hybrid peptide acting through GIP, glucagon and GLP-1 receptors exhibits weight reducing and anti-diabetic properties. *Biochemical Pharmacology* **8** 1655-1662.
- Black C, Donnelly P, McIntyre L, Royle P, Shepherd JJ & Thomas S 2009 Meglitinide analogues for type 2 diabetes mellitus (Review). *The Cochrane Collaboration* 1-52.

Bogacka I, Xie H, Bray GA & Smith SR 2004 The effect of Pioglitazone on peroxisome proliferator-activated receptor- γ target genes related to lipid storage in vivo. *Diabetes Care* **27** 1660–1667.

Bollag RJ, Zhong Q, Phillips P, Min L, Zhong L, Cameron R, Mulloy AL, Rasmussen H, Qin F, Ding KH & Isales CM 2000 Osteoblast-derived cells express functional glucose-dependent insulinotropic peptide receptors. *Endocrinology* **141** 1228-1235.

Bollag RJ, Zhong Q, Ding KH, Phillips P, Zhong L, Qin F, Cranford J, Mulloy AL, Cameron R & Isales CM 2001 Glucose-dependent insulinotropic peptide is an integrative hormone with osteotropic effects. *Molecular and Cellular Endocrinology* **177** 35-41.

Bonner-Weir S, Trent DF, Honey RN & Weir GC 1981 Responses of neonatal rat islets to streptozotocin: limited beta-cell regeneration and hyperglycemia. *Diabetes* **30** 64-69.

Bose AK, Mocanu MM, Carr RD, Brand CL & Yellon DM 2005 Glucagon-like peptide 1 can directly protect the heart ischemia/reperfusion injury. *Diabetes* **54** 146-151.

Bosi E, Camisasca RP, Collober C, Rochotte E & Garber AJ 2007 Effects of Vildagliptin on glucose control over 24 weeks in patients with type 2 diabetes inadequately controlled with metformin. *Diabetes Care* **30** 890-895.

Botolin S, Faugere M, Malluche H, Orth M, Meyer R & McCabe LR 2005 Increased bone adiposity and peroxisomal proliferator-activated receptor-2 expression in type 1 diabetic mice. *Endocrinology* **146** 3622-3631.

Botolin S & McCabe LR 2007 Bone loss and increased bone adiposity in spontaneous and pharmacologically induced diabetic mice. *Endocrinology* **148** 198-205.

Bouillon R 1991 Diabetic bone disease. *Calcified Tissue International* **49** 155-160.

Bouxsein ML, Boyd SK, Christiansen BA, Guldberg RE, Jepsen KJ & Müller R 2010 Guidelines for assessment of bone microstructure in rodents using microcomputed tomography. *Journal of Bone and Mineral Research* **25** 1468-1486.

Brownlee M 2001 Biochemistry and molecular cell biology of diabetic complications. *Nature Medicine* **414** 813-820.

Brubaker PL & Drucker DJ 2002 Structure-function of the glucagon receptor family of G protein-coupled receptors: the glucagon, GIP, GLP-1, and GLP-2 receptors. *Receptors and Channels* **8** 179-188.

Buse JB, Drucker DJ, Taylor KL, Kim T, Walsh B, Hu H, Wilhelm K, Trautmann M, Shen LZ & Porter LE 2010 DURATION-1: Exenatide once weekly produces sustained glycemic control and weight loss over 52 weeks. *Diabetes Care* **33** 1255-1261.

Cao JJ, Gregoire BR & Gao H 2009 High-fat diet decreases cancellous bone mass but has no effect on cortical bone mass in the tibia in mice. *Bone* **44** 1097-1104.

Ceccarelli E, Guarino EG, Merlotti D, Patti A, Gennari L, Nuti R & Dotta F 2013 Beyond glycemic control in diabetes mellitus : effects of incretin-based therapies on bone metabolism. *Frontiers in Endocrinology Bone Research* **4** 1-12.

Chachin M, Yamada M, Fujita A, Matsuoka T, Matsushita K & Kurachi Y 2003 Nateglinide, a D-phenylalanine derivative lacking either a sulfonylurea or benzamido moiety, specifically inhibits pancreatic beta-cell-type K_{ATP} channels. *The Journal of Pharmacology and Experimental Therapeutics* **304** 1025-1032.

Chao EC & Henry RR 2010 SGLT2 inhibition - a novel strategy for diabetes treatment. *Nature Reviews Drug Discovery* 1-9.

Chappard D, Basle MF, Legrand E & Audran M 2011 New laboratory tools in the assessment of bone quality. *Osteoporosis International* **22** 2225-2240.

Charbonnel B, Karasik A, Liu J, Wu M & Meininger G 2006 Efficacy and safety of the dipeptidyl peptidase-4 inhibitor Sitagliptin added to ongoing Metformin therapy in patients with type 2 diabetes inadequately controlled with Metformin alone. *Diabetes Care* **29** 2638-2643.

Chen H, Charlat O, Tartaglia LA, Woolf EA, Weng X, Ellis SJ, Lakey ND, Culpepper J, More KJ, Breitbart RE, Duyk GM, Tepper RI & Morgenstern JP 1996 Evidence That the Diabetes Gene Encodes the Leptin Receptor: Identification of a Mutation in the Leptin Receptor Gene in *db/db* Mice. *Cell* **84** 491-495.

Chen M, Bergman RN & Porte D 1988 Insulin resistance and beta cell dysfunction in aging : the importance of dietary carbohydrate. *Journal of Clinical Endocrinology & Metabolism* **67** 951-957.

Christensen M, Knop FK, Visboll T & Holst JJ 2011 Lixisenatide for type 2 diabetes Mellitus. *Expert Opinion in Investigational Drugs* **20** 549-557.

Coe LM, Zhang J & McCabe LR 2013 Both spontaneous *Ins2^{+/-}* and streptozotocin-induced type I diabetes cause bone loss in young mice. *Journal of Cellular Physiology* **228** 689-695.

Cohen MA, Ellis SM, Le Roux CW, Batterham RL, Park A, Patterson M, Frost GS, Ghatti MA & Bloom SR 2003 Oxyntomodulin suppresses appetite and reduces food intake in humans. *The Journal of Clinical Endocrinology & Metabolism* **88** 4696-4701.

Coleman DL 1978 Obese and diabetes : two mutant genes causing diabetes-obesity syndrome in mice. *Diabetologia* **14** 141-148.

Cornish J, Callon KE & Reid IR 1996 Insulin increases histomorphometric indices of bone formation *in vivo*. *Calcified Tissue International* **59** 492-495.

Cornish J, Callon K, Bava U, Lin C, Naot D, Hill B, Grey A, Broom N, Myers D, Nicholson G & Reid I 2002 Leptin directly regulates bone cell function *in vitro* and reduces bone fragility *in vivo*. *Journal of Endocrinology* **175** 405-415.

Culina S, Brezar V & Mallone R 2013 Mechanisms in endocrinology: Insulin and type 1 diabetes: immune connections. *European Journal of Endocrinology* **168** R19-R31.

Creutzfeldt W, Ebert R, Willms B, Frerichs H & Brown JC 1978 Gastric inhibitory polypeptide (GIP) and insulin in obesity: Increased response to stimulation and defective feedback control of serum levels. *Diabetologia* **14** 15-24.

Cummings SR, Bates D & Black DM 2002 Clinical use of bone densitometry: scientific review. *Scientific Review and Clinical Applications* **288** 1889-1897.

Dacquin R, Davey RA, Laplace C, Levasseur R, Morris HA, Goldring SR, Gebre-Medhin S, Galson DL, Zajac JD & Karsenty G 2004 Amylin inhibits bone resorption while the calcitonin receptor controls bone formation *in vivo*. *The Journal of Cell Biology* **164** 509-514.

Deacon CF, Knudsen LB, Madsen K, Wiberg FC, Jacobsen O & Holst JJ 1998 Dipeptidyl peptidase IV resistant analogues of glucagon-like peptide-1 which have extended metabolic stability and improved biological activity. *Diabetologia* **41** 271-278.

Deacon CF, Nauck MA, Meier J, Hucking K & Holst JJ 2000 Degradation of endogenous and exogenous gastric inhibitory polypeptide in healthy and in type 2 diabetic subjects as revealed using a new assay for the intact peptide. *Journal of Clinical Endocrinology & Metabolism* **8** 3575-3581.

Deacon CF, Danielsen P, Klarskov L, Olesen M & Holst JJ 2001 Dipeptidyl peptidase IV inhibition reduces the degradation and clearance of GIP and potentiates its insulintropic and antihyperglycemic effects in anesthetized pigs. *Diabetes* **50** 1588–1597.

Deacon CF, Wamberg S, Bie P, Hughes TE & Holst JJ 2002 Preservation of active incretin hormones by inhibition of dipeptidyl peptidase IV suppresses meal-induced incretin secretion in dogs. *Journal of Endocrinology* **172** 355–362.

Deacon CF, Plamboeck A, Rosenkilde MM de Heer J & Holst JJ 2006 GIP-(3– 42) does not antagonize insulintropic effects of GIP at physiological concentrations. *American Journal of Physiology - Endocrinology & Metabolism* **291** E468–E475.

DeFronzo RA, Ratner RE, Han J, Kim DD, Fineman MS & Baron AD 2005 Effects of Exenatide (Exendin-4) on glycemic control and weight over 30 weeks in Metformin-treated patients with type 2 diabetes. *Diabetes Care* **28** 1092–1100.

DeFronzo RA, Hissa MN, Garber AJ, Gross JL, Duan RY, Ravichandran S & Chen RS 2009 The efficacy and safety of Saxagliptin when added to Metformin therapy in patients with inadequately controlled type 2 diabetes with Metformin alone. *Diabetes Care* **32** 1649–1655.

DeFronzo RA, Tripathy D, Schwenke DC, Banerji MA, Bray GA, Buchanan TA, Clement SC, Henry RR, Hodis HN, Kitabchi AE, Mack WJ, Mudaliar S, Ratner RE, Williams K, Stentz FB, Musi N & Reaven PD 2011 Pioglitazone for diabetes prevention in impaired glucose tolerance. *The New England Journal of Medicine* **364** 1104–15.

Degn KB, Juhl CB, Sturis J, Jakobsen G, Brock B, Chandramouli V, Rungby J, Landau BR & Schmitz O 2004 One week's treatment with the long-acting glucagon-like peptide 1 derivative Liraglutide (NN2211) markedly improves 24-h glycemia and α - and beta-cell function and reduces endogenous glucose release in patients with type 2 diabetes. *Diabetes* **53** 1187–1194.

De Heer J, Rasmussen C, Coy DH & Holst JJ 2008 Glucagon-like peptide-1, but not glucose-dependent insulintropic peptide, inhibits glucagon secretion via somatostatin (receptor subtype 2) in the perfused rat pancreas. *Diabetologia* **51** 2263–2270.

De Liefde II, Van der Klift M, De Laet CEDH, Van Daela PLA, Hofman A & Pols HAP 2005 Bone mineral density and fracture risk in type-2 diabetes mellitus: the Rotterdam Study. *Osteoporos International* **16** 1713–1720.

Dempster DW, Compston JE, Drezner MK, Glorieux FH, Kanis JA, Malluche H, Meunier PJ, Ott SM, Recker RR & Parfitt AM 2013 Standardized nomenclature, symbols, and units for bone histomorphometry: A 2012 update of the report of the ASBMR Histomorphometry Nomenclature Committee. *Journal of Bone and Mineral Research* **28** 2-17.

Diabetes UK 2012 News [Online]
http://www.diabetes.org.uk/about_us/news_landing_page/nhs-spending-on-diabetes-to-reach-169-billion-by-2035/

Diabetes UK 2014 Diabetes complications [Online]
<http://www.diabetes.org.uk/guide-to-diabetes/complications/>

Ding K, Shi X, Zhong Q, Kang B, Xie D, Bollag WB, Bollag RJ, Hill W, Washington W, Mi Q, Insogna K, Chutkan N, Hamrick M & Isles CM 2008 Impact of glucose-dependent insulintropic peptide on age-induced bone loss. *Journal of Bone and Mineral Research* **23** 536-543.

Ding K, Zhong Q, Xie D, Chen H, Della-Fera MA, Bollag RJ, Bollag WB, Gujral R, Kang B, Sridhar S, Baile C, Curl W & Isles CM 2006 Effects of glucose-dependent insulintropic peptide on behavior. *Peptides* **27** 2750-2755.

Ding W & Gromada J 1997 Protein kinase A-dependent stimulation of exocytosis in mouse pancreatic β -cells by glucose-dependent insulintropic polypeptide. *Diabetes* **46** 615-621.

Dong XN, Qin A, Xu J & Wang X 2011 In situ accumulation of advanced glycation endproducts (AGEs) in bone matrix and its correlation with osteoclastic bone resorption. *Bone* **49** 174-183.

Dong JZ, Shen Y, Zhang J, Tsomaia N, Mierke DF & Taylor JE 2010 Discovery and characterization of Taspoglutide, a novel analogue of human glucagon-like peptide-1, engineered for sustained therapeutic activity in type 2 diabetes. *Diabetes, Obesity and Metabolism* **13** 19–25.

Drucker DJ, Philippe J, Mojsov S, Chick WL & Habener JF 1987 Glucagon-like peptide I stimulates insulin gene expression and increases cyclic AMP levels in a rat islet cell line. *Proceedings of the National Academy of Sciences* **84** 3434-3438.

Drucker DJ 2003 Enhancing incretin action for the treatment of type 2 diabetes. *Diabetes Care* **26** 2929-2940.

Drucker DJ & Nauck MA 2006 The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. *The Lancet* **368** 1696-1705.

Drucker DJ, Sherman SI, Gorelick FS, Bergenstal RM, Sherwin RS & Buse JB 2010 Incretin-based therapies for the treatment of type 2 diabetes: evaluation of the risks and benefits. *Diabetes Care* **33** 428-433.

Ducy P 2011 The role of osteocalcin in the endocrine cross-talk between bone remodeling and energy metabolism. *Diabetologia* **54** 1291-1297.

Ducy P, Amling M, Takeda S, Priemel M, Schilling AF, Beil FT, Shen J, Vinson C, Rueger JM & Karsenty G 2000 Leptin inhibits bone formation through a hypothalamic relay: a central control of bone mass. *Cell* **100** 197–207.

Duez H, Cariou B & Staels B 2012 DPP-4 inhibitors in the treatment of type 2 diabetes. *Biochemical Pharmacology* **83** 823-832.

Dunphy JL, Taylor RG & Fuller PJ 1998 Tissue distribution of rat glucagon receptor and GLP-1 receptor gene expression. *Molecular and Cellular Endocrinology* **141** 179–186.

During MJ, Cao L, Zuzga DS, Francis JS, Fitzsimons HL, Jiao X, Bland RJ, Klugmann M, Banks WA, Drucker DJ & Haile CH 2003 Glucagon-like peptide-1 is involved in learning and neuroprotection. *Nature Medicine* **9** 1173-1179.

Ealey KN, Fonseca D, Archer MC & Ward WE 2006 Bone abnormalities in adolescent leptin-deficient mice. *Regulatory Peptides* **136** 9-13.

Eastell R & Hannon RA 2008 Biomarkers of bone health and osteoporosis risk. *Proceedings of the Nutrition Society* **67** 157–162.

Ebert R, Unger H & Creutzfeldt W 1983 Preservation of incretin activity after removal of gastric inhibitory polypeptide (GIP) from rat gut extracts by immunoabsorption. *Diabetologia* **24** 449-454.

Ehses JA, Casilla VR, Doty T, Pospisilik JA, Winter KD, Demuth H, Pederson RA & McIntosh CHS 2003 Glucose-dependent insulinotropic polypeptide promotes beta-(INS-1) cell survival via cyclic adenosine monophosphate-mediated caspase-3 inhibition and regulation of p38 mitogen-activated protein kinase. *Endocrinology* **144** 4433-4445.

Eleazu CO, Eleazu KC, Chukwuma S & Essien UN 2013 Review of the mechanism of cell death resulting from streptozotocin challenge in experimental animals, its practical use and potential risk to humans. *Journal of Diabetes & Metabolic Disorders* **12** 60-66.

Eleftheriou F, Ahn JD, Takeda S, Starbuck M, Yang X, Liu X, Kondo H, Richards WG, Bannon TW, Noda M, Clement K, Vaisse C & Karsenty G 2005 Leptin regulation of bone resorption by the sympathetic nervous system and CART. *Nature* **434** 514-520.

Eng J, Kleinman WA, Singh L, Singh G & Raufman JP 1992 Isolation and characterization of exendin-4, an exendin-3 analogue, from *Heloderma suspectum* venom. Further evidence for an exendin receptor on dispersed acini from guinea pig pancreas. *Journal of Biological Chemistry* **267** 7402-7405.

Epstein S & LeRoith D 2008 Diabetes and fragility fractures — A burgeoning epidemic?. *Bone* **43** 3-6.

Erben RG & Glosmann M 2012 Histomorphometry in rodents. *Bone Research Protocols, Methods in Molecular Biology* **816** 279-303.

Ferron M, Wei J, Yoshizawa T, Fattore AD, DePinho RA, Teti A, Ducy P & Karsenty G 2010 Insulin signaling in osteoblasts integrates bone remodeling and energy metabolism. *Cell* **142** 296–308.

Filvaroff E, Erlebacher A, Ye J, Gitelman SE, Lotz J, Heilman M & Derynck R 1999 Inhibition of TGF-beta receptor signaling in osteoblasts leads to decreased bone remodeling and increased trabecular bone mass. *Development* **126** 4267-4279.

Fineman MS, Koda JE, Shen LZ, Strobel SA, Maggs DG, Weyer C & Koltermann OG 2002 The human amylin analog, pramlintide, corrects postprandial hyperglucagonemia in patients with type 1 diabetes. *Metabolism* **51** 636–641.

Fineman MS, Bicsak TA, Shen LZ, Taylor K, Gaines E, Varns A, Kim D & Baron AD 2003 Effect on glycemic control of Exenatide (synthetic Exendin-4) additive to existing metformin and/or sulfonylurea treatment in patients with type 2 Ddiabetes. *Diabetes Care* **26** 2370-2377.

Flatt PR, Bailey CJ, Kwasowski P, Swanson-Flatt SK & Marks V 1983 Abnormalities of GIP in spontaneous syndromes of obesity and diabetes in mice. *Diabetes* **32** 433-435.

Fonseca VA, Alvarado-Ruiz R, Raccach D, Boka G, Miossec P & Gerich JE 2012 Efficacy and safety of the once-daily GLP-1 receptor agonist Lixisenatide in monotherapy. *Diabetes Care* **35** 1225-1231.

Fonseca V, Schweizer A, Albrecht D, Baron MA, Chang I & Dejager S 2007 Addition of vildagliptin to insulin improves glycaemic control in type 2 diabetes. *Diabetologia* **50** 1148-1155.

Fowlkes JL, Bunn RC, Liu L, Wahl EC, Coleman HN, Cockrell GE, Perrien DS, Lumpkin Jr. CK & Thrailkill KM 2008 Runt-related transcription factor-2 (RUNX2) and RUNX2-related osteogenic genes are down-regulated throughout osteogenesis in type 1 diabetes mellitus. *Endocrinology* **149** 1697-1704.

Fratzl-Zelman N, Fratzl P, Hörandner H, Grabner B, Varga F, Ellinger A & Klaushofer K 1998 Matrix mineralization in MC3T3-E1 cell cultures initiated by beta-glycerophosphate pulse. *Bone* **23** 511-520.

Friedman JM & Halaas JL 1998 Leptin and the regulation of body weight in mammals. *Nature* **395** 763-770.

Fuhendorff J, Rorsman P, Kofod H, Brand CL, Rolin B, MacKay P, Shymko R & Carr RD 1998 Stimulation of insulin release by Repaglinide and Glibenclamide involves both common and distinct processes. *Diabetes* **47** 345-351.

Fulzele K & Clemens TL 2012 Novel functions for insulin in bone. *Bone* **50** 452-456.

Galluzzi F, Stagi S, Salti R, Toni S, Piscitelli E, Simonini G, Falcini F & Chiarelli F 2005 Osteoprotegerin serum levels in children with type 1 diabetes: a potential modulating role in bone status. *European Journal of Endocrinology* **153** 879-885.

Gault VA, Flatt PR, Bailey CJ, Harriott P, Greer B, Mooney MH & O'Harte FPM 2002 Enhanced cAMP generation and insulin-releasing potency of two novel

Tyr¹-modified enzyme-resistant forms of glucose-dependent insulintropic polypeptide is associated with significant antihyperglycaemic activity in spontaneous obesity-diabetes. *Biochemical Journal* **367** 913-920.

Gault VA, Flatt PR & O'Harte FPM 2003a Breakthroughs and views glucose-dependent insulintropic polypeptide analogues and their therapeutic potential for the treatment of obesity-diabetes. *Biochemical and Biophysical Research Communications* **308** 207–213.

Gault VA, O'Harte FPM, Harriott P, Mooney MH, Green BD & Flatt PF 2003b Effects of the novel (Pro³)GIP antagonist and exendin(9–39)amide on GIP- and GLP-1-induced cyclic AMP generation, insulin secretion and postprandial insulin release in obese diabetic (*ob/ob*) mice: evidence that GIP is the major physiological incretin. *Diabetologia* **46** 222-230.

Gault VA, Porter WD, Flatt PR & Holscher C 2010 Actions of Exendin-4 therapy on cognitive function and hippocampal synaptic plasticity in mice fed a high-fat diet. *International Journal of Obesity* **34** 1341-1344.

Gault VA, Lennox R & Flatt PR 2015 Sitagliptin, a dipeptidyl peptidase-4 inhibitor, improves recognition memory, oxidative stress and hippocampal neurogenesis and upregulates key genes involved in cognitive decline. *Diabetes, Obesity and Metabolism* **17** 403–413.

Giustina A, Mazziotti G & Canalis E 2008 Growth hormone, insulin-like growth factors and the skeleton. *Endocrine Reviews* **29** 535-559.

GlaxoSmithKline 2015 Tanzeum (Albiglutide) [Online] <http://us.gsk.com/en-us/media/press-kits/tanzeum-albiglutide/#full-us-prescribing-information>.

Göke R, Larsen PJ, Mikkelsen JD & Sheikh SP 1995 Distribution of GLP-1 Binding Sites in the Rat Brain: Evidence that Exendin-4 is a Ligand of Brain GLP-1 Binding Sites. *European Journal of Neuroscience* **7** 2294-2300.

Golub EE & Boesze-Battaglia K 2007 The role of alkaline phosphatase in mineralization. *Current Opinion in Orthopaedics* **18** 444-448.

Gomis R, Owens DR, Taskinen M, Del Prato S, Patel S, Pivovarov A, Schlosser A & Woerle H 2012 Long-term safety and efficacy of Linagliptin as monotherapy or in combination with other oral glucose-lowering agents in 2121 subjects with type 2 diabetes: up to 2 years exposure in 24-week phase III trials followed by a 78-week open-label extension. *International Journal of Clinical Practice* **66** 731-740.

Graham ML, Janecek JL, Kittredge JA, Hering BJ & Schuurman H 2011 The streptozotocin-induced diabetic nude mouse model: differences between animals from different sources. *Comparative Medicine* **61** 356-360.

Green BD & Flatt PR 2007 Incretin hormone mimetics and analogues in diabetes therapeutics. *Best Practice & Research Clinical Endocrinology & Metabolism* **21** 497-516.

Green BD, Flatt PR & Bailey CJ 2006 Dipeptidyl peptidase IV (DPP IV) inhibitors: a newly emerging drug class for the treatment of type 2 diabetes. *Diabetes and Vascular Disease Research* **3** 159-165.

Gregory CA, Gunn WG, Peister A & Prockop DJ 2004 An Alizarin red-based assay of mineralization by adherent cells in culture: comparison with cetylpyridinium chloride extraction. *Analytical Biochemistry* **329** 77-84.

Gremlich S, Porret A, Hani EH et al. (1995) Cloning, functional expression, and chromosomal localization of the human pancreatic islet glucose-dependent insulinotropic polypeptide receptor. *Diabetes* **44** 1202-1208.

Gromada J, Holst JJ & Roszman P 1998 Cellular regulation of islet hormone secretion by the incretin hormone glucagon-like peptide 1. *European Journal of Physiology* **435** 583-594.

Gulcelik NE, Bayraktar M, Caglar O, Alpaslan M & Karakaya J 2011 Mortality after hip fracture in diabetic patients. *Experimental and Clinical Endocrinology & Diabetes* **119** 414-418.

Halade GV, Rahman MM, Williams PJ & Fernandes G 2010 High fat diet-induced animal model of age-associated obesity and osteoporosis. *The Journal of Nutritional Biochemistry* **21** 1162-1169.

Hampson G, Evans C, Petitt RJ, Evans WD, Woodhead SJ, Peters JR & Ralston SH 1998 Bone mineral density, collagen type 1 $\alpha 1$ genotypes and bone turnover in premenopausal women with diabetes mellitus. *Diabetologia* **41** 1314-1320.

Hamrick MW, Pennington C, Newton D, Xie D & Isles C 2004 Leptin deficiency produces contrasting phenotypes in bones of the limb and spine. *Bone* **34** 376-383.

Henriksen DB, Alexandersen P, Bjarnason NH, Vilsbøll T, Hartmann B, Henriksen EE, Byrjalsen I, Krarup T, Holst JJ & Christiansen C 2003 Role of gastrointestinal hormones in postprandial reduction of bone resorption. *Journal of Bone and Mineral Research* **18** 2180-2189.

Herman GA, Stevens C, Van Dyck K, Bergman A, Yi B, De Smet M, Snyder K, Hilliard D, Tanen M, Tanaka W, Wang AQ, Zeng W, Musson D, Winchell G, Davies MJ, Ramael S, Gottesdiener KM & Wagner JA 2005 Pharmacokinetics and pharmacodynamics of Sitagliptin, an inhibitor of dipeptidyl peptidase IV, in healthy subjects : results from two randomized, double-blind, placebo-controlled studies with single oral doses. *Clinical Pharmacology and Therapeutics* **78** 675-688.

Hermann LS, Scherstén B, Bitzén P, Kjellström T, Lindgärde F & Melander A 1994 Therapeutic comparison of metformin and sulfonylurea, alone and in various combinations: A double-blind controlled study. *Diabetes Care* **17** 1100-1109.

Herrmann C, Göke R, Richter G, Fehmann HC, Arnold R & Göke B 1995 Glucagon-like peptide-1 and glucose-dependent insulin-releasing polypeptide plasma levels in response to nutrients. *Digestion* **56** 117-26.

Hernandez CJ & Keaveny TM 2006 A biomechanical perspective on bone quality. *Bone* **39** 1173-1181.

Hessle L, Johnson KA, Anderson HC, Narisawa S, Sali A, Goding JW, Terkeltaub R & Millan JL 2002 Tissue-nonspecific alkaline phosphatase and plasma cell membrane glycoprotein-1 are central antagonistic regulators of bone mineralization. *Proceedings of the National Academy of Sciences of the United States of America* **99** 9445-9449.

Hinke SA, Gelling RW, Pederson RA, Manhart S, Nian C, Demuth H & McIntosh CHS 2002 Dipeptidyl peptidase IV-resistant [d-Ala²]glucose-dependent insulinotropic polypeptide (GIP) improves glucose tolerance in normal and obese diabetic rats. *Diabetes* **51** 652-661.

Hofbauer LC, Brueck CC, Singh SK & Dobnig H 2007 Osteoporosis in patients with diabetes mellitus. *Journal of Bone and Mineral Research* **22** 1317-1328.

Holecki M & Wiecek A 2010 Relationship between body fat mass and bone metabolism. *Polish Archives of Internal Medicine* **120** 361-167.

Hollander PL, Li J, Frederick R, Allen E & Chen R 2011 Safety and efficacy of Saxagliptin added to Thiazolidinedione over 76 weeks in patients with type 2 diabetes mellitus. *Diabetes and Vascular Disease Research* **8** 125-135.

Holloway WR, Collier FM, Aitken CJ, Myers DE, Hodge JM, Malakellis M, Gough TJ, Collier GR & Nicholson GC 2002 Leptin inhibits osteoclast generation. *Journal of Bone and Mineral Research* **17** 200-209.

Holscher C 2012 Potential role of glucagon-like peptide-1 (GLP-1) in neuroprotection. *CNS Drugs* **2012** 871-882.

Holst JJ & Deacon CF 1998 Inhibition of the activity of dipeptidyl-peptidase IV as a treatment for type 2 diabetes. *Diabetes* **47** 1663-1670.

Holst JJ, Christensen M, Lund A, de Heer J, Svendsen B, Kielgast U & Knop FK 2011 Regulation of glucagon secretion by incretins. *Diabetes, Obesity and Metabolism* **13** 89–94.

Horcajada-Molteni M, Chanteranne B, Lebecque P, Davicco M, Coxam V, Young A & Barlet J 2001 Amylin and bone metabolism in streptozotocin-induced diabetic rats. *Journal of Bone and Mineral Research* **16** 958-965.

Horowitz M, Rayner CK & Jones KL 2013 Mechanisms and clinical efficacy of Lixisenatide for the management of type 2 diabetes. *Advances in Therapy* **30** 81–101.

Hu FB 2011 Globalization of diabetes : the role of diet, lifestyle and gene. *Diabetes Care* **34** 1249-1257.

Hu S, Wang S, Fanelli B, Bell PA, Dunning BE, Geisse S, Schmitz R & Boettcher BR 2000 Pancreatic beta-cells KATP channel activity and membrane-binding studies with nateglinide : a comparison with sulphonylureas and repaglinide. *The Journal of Pharmacology and Experimental Therapeutics* **293** 444-452.

International Diabetes Federation 2014a About diabetes [Online]
<http://www.idf.org/about-diabetes>

International Diabetes Federation 2014b IDF diabetes atlas [Online]
<http://www.idf.org/diabetesatlas/update-2014>

Inzucchi SE 2002 Oral antihyperglycaemic therapy for type 2 diabetes scientific review. *The Journal of the American Medical Association* **287** 360-372.

Inzucchi SE, Bergenstal RM, Buse JB, Diamant M, Ferrannini E, Nauck M, Peters AL, Tsapas A, Wender R & Matthews DR 2012 Management of

hyperglycaemia in type 2 diabetes: a patient-centered approach. Position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* **35** 1364-1379.

Ionova-Martin SS, Do SH, Barth HD, Szadkowska M, Porter AE, Ager III JW, Ager Jr JW, Alliston T, Vaisse C & Ritchie RO 2010 Reduced size-independent mechanical properties of cortical bone in high-fat diet-induced obesity. *Bone* **46** 217-225.

Irwin N, Green BD, Mooney MH, Greer B, Harriott P, Bailey CJ, Gault VA, O'Harte FPM & Flatt PR 2005a A novel, long-acting agonist of glucose-dependent insulintropic polypeptide suitable for once-daily administration in type 2 diabetes. *Journal of Pharmacology and Experimental Therapeutics* **314** 1187-1194.

Irwin N, Green BD, Gault VA, Greer B, Harriott P, Bailey CJ, Flatt PR & O'Harte FPM 2005b Degradation, insulin secretion and antihyperglycemic actions of two palmitate-derivitized N-terminal pyroglutamyl analogues of glucose-dependent insulintropic polypeptide. *Journal of Medicinal Chemistry* **48** 1244-1250.

Irwin N, O'Harte FPM, Gault VA, Green BD, Greer B, Harriott P, Bailey CJ & Flatt PR 2006a GIP(Lys¹⁶PAL) and GIP(Lys³⁷PAL): novel long-acting acylated analogues of glucose-dependent insulintropic polypeptide with improved antidiabetic potential. *Journal of Medicinal Chemistry* **49** 1047-1054.

Irwin N, Clarke GC, Green BD, Greer B, Harriott P, Gault VA, O'Harte FPM & Flatt PR 2006b Evaluation of the antidiabetic activity of DPP IV resistant N-terminally modified versus mid-chain acylated analogues of glucose-dependent insulintropic polypeptide. *Biochemical Pharmacology* **72** 719-728.

Irwin N & Flatt PR 2009 Therapeutic potential for GIP receptor agonists and antagonists. *Best Practice & Research Clinical Endocrinology & Metabolism* **23** 499–512.

Irwin N, Gault V & Flatt PR 2010 Therapeutic potential of the original incretin hormone glucose-dependent insulintropic polypeptide : diabeted, obesity, osteoporosis and Alzheimer's disease?. *Expert Opinion on Investigational Drug* **19** 1039-1048.

Janghorbani M, Dam RMV, Willett WC & Hu FB 2007 Systematic review of type 1 and type 2 diabetes mellitus and risk of fracture. *American Journal of Epidemiology* **166** 495-505.

Janssens K, Dijke PT, Janssens S & Hul WV 2005 Transforming growth factor-beta1 to the bone. *Endocrine Reviews* **6** 743-774.

Jehle P, Jehle D, Mohan S & Bohm B 1998 Serum levels of insulin-like growth factor system components and relationship to bone metabolism in Type 1 and Type 2 diabetes mellitus patients. *Journal of Endocrinology* **159** 297-306.

Johnson ML, Harnish K, Nusse R & Hul WV 2004 Review LRP5 and Wnt signaling: a union made for bone. *Journal of Bone and Mineral Research* **19** 1749-1757.

Justesen J, Stenderup K, Ebbesen EN, Mosekilde L, Steiniche T & Kassem M 2001 Adipocyte tissue volume in bone marrow is increased with aging and in patients with osteoporosis. *Biogerontology* **2** 165-171.

Kahn SE, Haffner SM, Heise MA, Herman WH, Holman RR, Jones, NP, Kravitz BG, Lachin JM, O'Neill MC, Zinman B & Viberti G 2006 Glycemic durability of Rosiglitazone, Metformin or Glyburide monotherapy. *The New England Journal of Medicine* **355** 2427-2443.

Kahn SE, Hull RL & Utzschneider KM 2006 Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature* **444** 840–846.

Kaplan AM & Vigna SR 1994 Gastric inhibitory polypeptide (GIP) binding sites in rat brain. *Peptides* **15** 297-302.

Karasik A, Aschner P, Katzeff H, Davies MJ & Stein PP 2008 Sitagliptin, a DPP-IV inhibitor for the treatment of patients with type 2 diabetes: a review of recent clinical trials. *Current Medical Research and Opinion* **24** 489-496.

Kendall DM, Riddle MC, Rosenstock J, Zhuang D, Kim DD, Fineman MS & Baron AD 2005 Effects of Exenatide (Exendin-4) on glycemic control over 30 weeks in patients with type 2 diabetes treated with Metformin and a Sulfonylurea. *Diabetes Care* **28** 1083-1091.

Khosla S 2001 Minireview: The OPG/RANKL/RANK system. *Endocrinology* **142** 5050–5055.

Kieffer TJ, McIntosh CHS & Pederson RA 1995 Degradation of glucose-dependent insulintropic polypeptide and truncated glucagon-like peptide 1 *in vitro* and *in vivo* by dipeptidyl peptidase IV. *Endocrinology* **136** 3585-3596.

Knudsen LB, Nielsen PF, Huusfeldt PO, Johansen NL, Madsen K, Pedersen FZ, Thøgersen H, Wilken M & Agersø H 2000 Potent derivatives of glucagon-like peptide-1 with pharmacokinetics properties suitable for once daily administration. *Journal of Medicinal Chemistry* **43** 1664-1669.

Koltermann OG, Buse JB, Fineman MS, Gaines E, Heintz S, Bicsak TA, Taylor K, Aisporna M, Wang Y & Baron AD 2003 Synthetic exendin-4 (exenatide) significantly reduces postprandial and fasting plasma glucose in subjects with type 2 diabetes. *The Journal of Clinical Endocrinology & Metabolism* **88** 3082-3089.

Kostenuik PJ 2005 Osteoprotegerin and RANKL regulate bone resorption, density, geometry and strength. *Current Opinion in Pharmacology* **5** 618-625.

Krentz AJ & Bailey CJ 2005 Oral antidiabetic agents current role in type 2 diabetes mellitus. *Drugs* **65** 385-411.

Kume S, Kato S, Yamagishi S, Inagaki Y, Ueda S, Arima N, Okawa T, Kojiro M & Nagata K 2005 Advanced glycation end-products attenuate human mesenchymal stem cells and prevent cognate differentiation into adipose tissue, cartilage and bone. *Journal of Bone and Mineral Research* **20** 1647-1658.

Kyle KA, Willett TL, Baggio LL, Drucker DJ & Grynps MD 2011 Differential Effects of PPAR-gamma activation versus chemical or genetic reduction of DPP-4 activity on bone quality in mice. *Endocrinology* **152** 457-467.

Laharrague P, Larrouy D, Fontanilles A, Truel N, Campfield A, Tenenbaum R, Galitzky J, Corberand JX, Penicaud L & Casteilla L 1998 High expression of leptin by human bone marrow adipocytes in primary culture. *The FASEB Journal* **12** 747-752.

Lamont BJ & Drucker DJ 2008 Differential antidiabetic efficacy of incretin agonists versus DPP-4 inhibition in high fat-fed mice. *Diabetes* **57** 190-198.

Lamothe B, Webster WK, Gopinathan A, Besse A, Campos AD & Darnay BG 2007 TRAF6 ubiquitin ligase is essential for RANKL signaling and osteoclast differentiation. *Biochemical and Biophysical Research Communications* **359** 1044-1049.

Larsson H, Holst JJ & Ahren B 1997 Glucagon-like peptide-1 reduces hepatic glucose production indirectly through insulin and glucagon in human. *Acta Physiologica Scandinavica* **160** 413-422.

Leahy JL 2005 Pathogenesis of Type 2 Diabetes Mellitus. *Archives of Medical Research* **36** 197-209.

Lecka-Czernik B, Moerman EJ, Grant DF, Lehmann JM, Manolagas SC & Jilka RL 2002 Divergent effects of selective peroxisome proliferator-activated receptor- γ 2 ligands on adipocyte *versus* osteoblast differentiation. *Endocrinology* **143** 2376-2384.

Lee AJ, Hodges S & Eastell R 2000 Measurement of osteocalcin. *Annals of Clinical Biochemistry* **37** 432-446.

Lee KS, Hong SH & Bae SC 2002 Both the Smad and p38 MAPK pathways play a crucial role in Runx2 expression following induction by transforming growth factor-beta and bone morphogenetic protein. *Oncogene* **21** 7156-7163.

Lee NK, Sowa H, Hinoi E, Ferron M, Ahn JD, Confavreux C, Dacquin R, Mee PJ, McKee MD, Jung DY, Zhang Z, Kim JK, Mauvais-Jarvis F, Ducy P & Karsenty G 2007 Endocrine regulation of energy metabolism by the skeleton. *Cell* **130** 456-469.

Lenzen S 2008 The mechanisms of alloxan- and streptozotocin-induced diabetes. *Diabetologia* **51** 216-226.

Li Y, Hansotia T, Yusta B, Ris F, Halban PA & Drucker DJ 2003 Glucagon-like peptide-1 receptor signaling modulates beta-cell apoptosis. *The Journal of Biological Chemistry* **278** 471-478.

Libouban H, Blouin S, Moreau M, Baslé MF, Audran M & Chappard D 2008 Effects of risedronate in a rat model of osteopenia due to orchidectomy and disuse: densitometric, histomorphometric and microtomographic studies. *Micron* **39** 998-1007.

Licata A 2009 Bone density vs bone quality : What's a clinician to do?. *Cleveland Clinic Journal of Medicine* **76** 331-336.

- List JF, Woo V, Morales E, Tang W & Fiedorek FT 2009 Sodium-glucose cotransport inhibition with Dapagliflozin in type 2 diabetes. *Diabetes Care* **32** 650–657.
- Lovshin, JA & Drucker, DJ 2009 Incretin-based therapies for type 2 diabetes mellitus. *Nature Reviews Endocrinology* **5** 262-269.
- Lu W, Juang J, Hsu BR & Huang H 1998 Effects of High or Low Dose of Streptozocin on Pancreatic Islets in C57BL/6 and C.B17-SCID Mice 1. *Transplantation Proceedings* **30** 609-610.
- Luna R, Garcia-Mayor RV, Lage M, Andrade MA, Barreiro J, Pombo M, Dieguez C & Casanueva FF 1999 High serum leptin levels in children with type 1 diabetes mellitus: contribution of age, BMI, pubertal development and metabolic status. *Clinical Endocrinology* **51** 603-610.
- Lund A, Knop FK & Visboll T 2011 Emerging GLP-1 receptor agonists. Expert Opinion in Emerging Drugs **16** 607-618.
- Lund A, Knop F & Visboll T 2014 Glucagon-like peptide-1 receptor agonists for the treatment of type 2 diabetes: differences and similarities. *European Journal of Internal Medicine* <http://dx.doi.org/10.1016/j.ejim.2014.03.005>.
- Mabilleau G, Mieczkowska A, Irwin N, Flatt PR & Chappard D 2013 Optimal bone mechanical and material properties require a functional glucagon-like peptide-1 receptor. *Journal of Endocrinology* **219** 59-68.
- Mabilleau G, Mieczkowska A, Irwin N, Simon Y, Audran M, Flatt PR & Chappard D 2014 Beneficial effects of a N-terminally modified GIP agonist on tissue-level bone material properties. *Bone* **63** 61-68.
- Mabilleau G, Mieczkowska A, Libouban H, Simon Y, Audran M & Chappard D 2015 Comparison between quantitative X-ray imaging, dual energy X-ray absorptiometry and microCT in the assessment of bone mineral density in

disuse-induced bone loss. *Journal of Musculoskeletal and Neuronal Interactions* **15** 42-52.

Mahoney DJ, Mikecz K, Ali T, Mabileau G, Benayahu D, Plaas A, Milner CM, Day AJ & Sabokbar A 2008 TSG-6 regulates bone remodeling through inhibition of osteoblastogenesis and osteoclast activation. *The Journal of Biological Chemistry* **283** 25952-25962.

Maida A, Hansotia T, Longuet C, Seino Y & Drucker DJ 2009 Differential importance of glucose-dependent insulintropic polypeptide vs glucagon-like peptide 1 receptor signaling for beta cell survival in mice. *Gastroenterology* **137** 2146–2157.

Mathers CD & Loncar D 2006 Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Medicine* **3** 2011-2030.

Matsunuma A, Kawane T, Maeda T, Hamada S & Horiuchi N 2004 Leptin corrects increased gene expression of renal 25-hydroxyvitamin D₃-1 α -hydrolase and -24-hydrolase in leptin-deficient, *ob/ob* mice. *Endocrinology* **145** 1367-1375.

Matsuo K & Irie N 2008 Osteoclast–osteoblast communication. *Archives of Biochemistry and Biophysics* **473** 201-209.

Mayo KE, Miller LJ, Bataille D, Dalle S, Goke B, Thorens B & Drucker DJ 2003 The glucagon receptor family. *Pharmacological Reviews* **55** 167-194.

McCabe LR 2007 Understanding the pathology and mechanisms of type I diabetic bone loss. *Journal of Cellular Biochemistry* **102** 1343-1357.

McCall M & Shapiro AMJ 2012 Update on islet transplantation. *Cold Spring Harbor Perspectives in Medicine* **2** 1-16.

- McCarthy AD, Etcheverry SB, Bruzzone L & Cortizo AM 1997 Effects of advanced glycation end-products on the proliferation and differentiation of osteoblast-like cells. *Molecular and Cellular Biochemistry* **170** 43-51.
- McIntosh CHS, Widenmaier S & Kim S 2009 Glucose-dependent insulintropic polypeptide (Gastric Inhibitory Polypeptide; GIP). *Vitamins and Hormones* **80** 409-471.
- Meier JJ, Nauck MA, Kranz D, Holst JJ, Deacon CF, Gaeckler D, Schmidt WE & Gallwitz B 2004 Secretion, degradation and elimination of glucagon-like peptide 1 and gastric inhibitory polypeptide in patients with chronic renal insufficiency and healthy control subjects. *Diabetes* **53** 654-662.
- Mentlein R, Gallwitz B & Schmidt WE 1993 Dipeptidyl-peptidase IV hydrolyses gastric inhibitory polypeptide, glucagon-like peptide-1(7-36)amide, peptide histidine methionine and is responsible for their degradation in human serum. *European Journal of Biochemistry* **214** 829-835.
- Merchenthaler I, Lane M & Shughrue P 1999 Distribution of pre-pro-glucagon and glucagon-like peptide-1 receptor messenger RNAs in the rat central nervous system. *The Journal of Comparative Neurology* **403** 261-80.
- Miazgowski T & Czekalski S 1998 A 2-year Follow-up study on bone mineral density and markers of bone turnover in patients with long-standing insulin-dependent diabetes mellitus. *Osteoporosis International* **8** 399-403.
- Mieczkowska A, Irwin N, Flatt PR, Chappard D & Mabileau G 2013 Glucose-dependent insulintropic polypeptide (GIP) receptor deletion leads to reduced bone strength and quality. *Bone* **56** 337-342.
- Miyata T, Notoya K, Yoshida K, Horie K, Maeda K, Kurokawa K & Taketomi S 1997 Advanced glycation end products enhance osteoclast-induced bone resorption in cultured mouse unfractionated bone cells and in rats implanted

subcutaneously with devitalized bone particles. *Journal of the American Society of Nephrology* **8** 260-270.

Moseley KF 2012 Type 2 diabetes and bone fractures. *Current Opinion in Endocrinology, Diabetes and Obesity* **19** 128-135.

Mojsov S, Weir GC & Habener JF 1983 Insulinotropin: glucagon-like peptide I (7-37) co-encoded in the glucagon gene is a potent stimulator of insulin release in the perfused rat pancreas. *Journal of Clinical Investigations* **79** 616-619.

Motyl K & McCabe LR 2009 Streptozotocin, type 1 diabetes severity and bone. *Biological Procedures Online* **11** 296-315.

Mu J, Woods J, Zhou Y, Roy RS, Li Z, Zycband E, Feng Y, Zhu L, Li C, Howard AD, Moller DE, Thornberry NA & Zhang BB 2006 Chronic Inhibition of Dipeptidyl Peptidase-4 With a Sitagliptin Analog Preserves Pancreatic β -Cell Mass and Function in a Rodent Model of Type 2 Diabetes. *Diabetes* **55** 1695-1704.

Nakamura M, Udagawa N, Matsuura S, Mogi M, Nakamura H, Horiuchi H, Saito N, Hiraoka BY, Kobayashi Y, Takaoka K, Ozawa H, Miyazawa H & Takahashi N 2003 Osteoprotegerin regulates bone formation through a coupling mechanism with bone resorption. *Endocrinology* **144** 5441-5449.

Natali A & Ferrannini E 2006 Effects of metformin and thiazolidinediones on suppression of hepatic glucose production and stimulation of glucose uptake in type 2 diabetes: a systematic review. *Diabetologia* **49** 434-441.

Nauck MA, Homberger E, Siegel BG, Allen RC, Eaton RP, Ebert R & Creutzfeldt W 1986 Incretin effects of increasing glucose loads in man calculated from venous insulin and C-peptide responses. *Journal of Clinical Endocrinology & Metabolism* **63** 492-498.

Nauck MA, Vilsbøll T, Gallwitz B, Garber A & Madsbad S 2009 Incretin-based therapies: viewpoints on the way to consensus. *Diabetes Care* **32** S223-S231.

Nauck MA, Vardarli I, Deacon CF, Holst JJ & Meier JJ 2011 Secretion of glucagon-like peptide-1 (GLP-1) in type 2 diabetes : what is up, what is down?. *Diabetologia* **54** 10-18.

Neumiller JJ & Campbell RK 2009 Liraglutide: a once-daily incretin mimetic for the treatment of type 2 diabetes mellitus. *The Annals of Pharmacotherapy* **43** 1433-1444.

Neumiller JJ 2012 Incretin pharmacology: a review of the incretin effect and current incretin-based therapies. *Cardiovascular & Hematological Agents in Medicinal Chemistry* **10** 276-288.

Nielsen LL, Young AA & Parkes DG 2004 Pharmacology of exenatide (synthetic exendin-4): a potential therapeutic for improved glycemic control of type 2 diabetes. *Regulatory Peptides* **117** 77-88.

Nikolaidis LA, Elahi D, Shen Y & Shannon RP 2004 Active metabolite of GLP-1 mediates myocardial glucose uptake and improves left ventricular performance in conscious dogs with dilated cardiomyopathy. *The American Journal of Physiology - Heart and Circulatory Physiology* **289** H2401-H2408.

Nuche-Berenguer B, Moreno P, Esbrit P, Dapia S, Caeiro JR, Cancelas J, Haro-Mora JJ & Villanueva-Peñacarrillo ML 2009 Effect of GLP-1 treatment on bone turnover in normal, type 2 diabetic, and insulin-resistant states. *Calcified Tissue International* **84** 453-461.

Nuche-Berenguer B, Portal-Núñez S, Moreno P, González N, Acitores A, López-Herradón A, Esbrit P, Valverde I & Villanueva-Peñacarrillo ML 2010a Presence of a functional receptor for GLP-1 in osteoblastic cells, independent of the cAMP-linked GLP-1 receptor. *Journal of Cellular Physiology* **225** 585-592.

Nuche-Berenguer B, Moreno P, Portal-Nuñez S, Dapía S, Esbrit P & Villanueva-Peñacarrillo ML 2010b Exendin-4 exerts osteogenic actions in insulin-resistant and type 2 diabetic states. *Regulatory Peptides* **159** 61-66.

Nyberg J, Jacobsson C, Anderson MF & Eriksson PS 2007 Immunohistochemical distribution of glucose-dependent insulinotropic polypeptide in the adult rat brain. *Journal of Neuroscience Research* **85** 2099-2119.

Nyholm B, Orskov L, Hove KY, Gravholt CH, Møller N, Alberti KGMM, Moyses C, Kolterman O & Schmitz O The amylin analog Pramlintide improves glycemic control and reduces postprandial glucagon concentrations in patients with type 1 diabetes mellitus. *Metabolism* **48** 935-941.

Nyman JS, Even JL, Jo C, Herbert EG, Murry MR, Cockrell GE, Wahl EC, Bunn RC, Lumpkin Jr CK, Fowlkes JL & Thrailkill KM 2011 Increasing duration of type 1 diabetes perturbs the strength-structure relationship and increases brittleness of bone. *Bone* **48** 733-740.

O'Harte FP, Mooney MH, Kelly CMN & Flatt PR 2000 Improved glycaemic control in obese diabetic ob/ob mice using N-terminally modified gastric inhibitory polypeptide. *The Journal of Endocrinology* **165** 639-648.

O'Harte FPM, Gault VA, Parker JC, Harriott P, Mooney MH, Bailey CJ & Flatt PR 2002 Improved stability, insulin-releasing activity and antidiabetic potential of two novel N-terminal analogues of gastric inhibitory polypeptide: N-acetyl-GIP and pGlu-GIP. *Diabetologia* **45** 1281-1291.

Oliver WC & Pharr GM 1992 An improved technique for determining hardness and elastic modulus using load and displacement sensing indentation experiments. *Journal of Materials Research* **7** 1564-1583.

Omar B, Banke E, Guiguis E, Åkesson L, Manganiello V, Lyssenko V, Groop L, Gomez MF & Degerman E 2012 Regulation of the pro-inflammatory cytokine osteopontin by GIP in adipocytes - a role for the transcription factor NFAT and

phosphodiesterase 3B. *Biochemical and Biophysical Research Communications* **425** 812–817.

Orriss IR, Taylor SEB & Arnett TR 2012 Rat osteoblasts cultures. *Bone Research Protocols, Methods in Molecular Biology* **816** 31-41.

Orskov C, Wettergren A & Holst JJ 1993 Biological effects and metabolic rates of glucagon-like peptide-1 (7-36)amide and glucagon-like peptide-1(7-37) in healthy subjects are indistinguishable. *Diabetes* **42** 658-661.

Orskov C, Rabenhøj L, Wettergren A, Kofod H & Holst JJ 1994 Tissue and plasma concentrations of amidated and glycine-extended glucagon-like peptide-1 in humans. *Diabetes* **43** 535-539.

Pamir N, Lynn FC, Buchan AMJ, Ehses J, Hinke SA, Pospisilik JA, Miyawaki K, Yamada Y, Seino Y, McIntosh CHS & Pederson RA 2003 Glucose-dependent insulinotropic polypeptide receptor null mice exhibit compensatory changes in the enteroinsular axis. *American Journal of Physiology - Endocrinology and Metabolism* **284** E931-E939.

Parkes D, Jodka C, Smith P, Nayak S, Rinehart L, Gingerich R, Chen K & Young A 2001 Pharmacokinetic actions of exendin-4 in the rat: Comparison with glucagon-like peptide-1. *Drug Development Research* **53** 260-267.

Paschalis EP 2012 Fourier transform infrared imaging of bone. *Bone Research Protocols, Methods in Molecular Biology* **816** 517-525.

Pasco JA, Henry MJ, Kotowicz MA, Collier GR, Ball MJ, Ugoni AM & Nicholson GC 2001 Serum leptin levels are associated with bone mass in nonobese women. *The Journal of Clinical Endocrinology & Metabolism* **86** 1884-1887.

Patsch JM, Kiefer FW, Varga P, Pail P, Rauner M, Stupphann D, Resch H, Moser D, Zysset PK, Stulnig TM & Pietschmann P 2011 Increased bone resorption and

impaired bone microarchitecture in short-term and extended high-fat diet-induced obesity. *Metabolism* **60** 243-249.

Patterson CC, Dahlquist GG, Gyürüs E, Green A & Soltész G 2009 Incidence trends for childhood type 1 diabetes in Europe during 1989–2003 and predicted new cases 2005–20: a multicentre prospective registration study. *The Lancet* **373** 2027-2033.

Patterson CC, Gyurus E, Rosenbauer J, Cinek O, Neu A, Schober E, Parslow RC, Joner G, Svensson J, Castell C, Bingley PJ, Schoenle E, Jarosz-Chobot P, Urbonaite B, Rothe U, Krzisnik C, Ionescu-Tirgoviste C, Weets I, Kocova M, Stipancic G, Samardzic M, de Beaufort CE, Green A, Dahlquist GG & Soltesz G 2012 Trends in childhood type 1 diabetes incidence in Europe during 1989-2008: evidence of non-uniformity over time in rates of increase. *Diabetologia* **55** 2142-2147.

Pautke C, Schieker M, Tischler T, Kolk A, Neth P, Mutschler W & Milz S 2004 Characterization of osteosarcoma cell lines MG-63, Saos-2 and U-2 OS in comparison to human osteoblasts. *Anticancer Research* **24** 3743-3748.

Pederson RA, Schubert HE & Brown JC 1975 Gastric inhibitory polypeptide: its physiologic release and insulinotropic action in the dog. *Diabetes* **24** 1050-1056.

Perry TA, Haughey NJ, Mattson MP, Egan JM & Greig NH 2002 Protection and reversal of excitotoxic neuronal damage by glucagon-like peptide-1 and Exendin-4. *The Journal of Pharmacology and Experimental Therapeutics* **302** 881-888.

Perry TA, Lahiri DK, Sambamurti K, Chen D, Mark P, Mattson MP, Egan JM & Greig NH 2003 Glucagon-like peptide-1 decreases endogenous amyloid- β peptide (A β) levels and protects hippocampal neurons from death induced by A β and iron. *Journal of Neuroscience Research* **72** 603– 612.

Peterson GE & Pollom RD 2010 Liraglutide in clinical practice: dosing, safety and efficacy. *The International Journal of Clinical Practice* **64** 35-43.

Peterson KF & Sullivan JT 2001 Effects of a glucagon receptor antagonist (Bay 27-9955) on glucagon-stimulated glucose production in humans. *Diabetologia* **44** 2018-2024.

Petit MA, Paudel ML, Taylor BC, Hughes JM, Strotmeyer ES, Schwartz AV, Cauley JA, Zmuda JM, Hoffman AR & Ensrud KE 2010 Bone mass and strength in older men with type 2 diabetes: the osteoporotic fractures in men study. *Journal of Bone and Mineral Research* **25** 285–291.

Pfeiffer AFH 2008 Oral hypoglycemic agents: Sulfonylureas and Meglitinides. *Type 2 Diabetes Principles and Practice Second Edition* 97-105.

Pinkse GGM, Tysma OHM, Bergen CAM, Kester MGD, Ossendorp F, van Veelen PA, Keymeulen B, Pipeleers D, Drijfhout JW & Roep BO 2005 Autoreactive CD8 T cells associated with β cell destruction in type 1 diabetes. *Proceedings of the National Academy of Sciences of the United States of America* **102** 18425-18430.

Pocai A 2012 Unraveling oxyntomodulin, GLP1's enigmatic brother. *Journal of Endocrinology* **215** 335–346.

Pocai A 2014 Action and therapeutic potential of oxyntomodulin. *Molecular Metabolism* **3** 241-251.

Porter DW, Kerr BD, Flatt PR, Holscher C & Gault VA 2010 Four weeks administration of Liraglutide improves memory and learning as well as glycaemic control in mice with high fat dietary-induced obesity and insulin resistance. *Diabetes, Obesity and Metabolism* **12** 891-899.

Porter DW, Irwin N, Flatt PR, Hölscher C & Gault VA 2011 Prolonged GIP receptor activation improves cognitive function, hippocampal synaptic plasticity and

glucose homeostasis in high-fat fed mice. *European Journal of Pharmacology* **650** 688-693.

Pospisilik JA, Martin J, Doty T, Ehses JA, Pamir N, Lynn FC, Piteau S, Demuth H, McIntosh CHS & Pederson RA 2003 Dipeptidyl Peptidase IV Inhibitor Treatment Stimulates β -Cell Survival and Islet Neogenesis in Streptozotocin-Induced Diabetic Rats. *Diabetes* **52** 741-750.

Pratley RE, Jauffret-Kamel S, Galbreath E & Holmes D 2006 Twelve-week monotherapy with the DPP-4 inhibitor vildagliptin improves glycaemic control in subjects with type 2 diabetes. *Hormone and Metabolic Research* **387** 423-428.

Preitner F, Ibberson M, Franklin I, Binnert C, Pende M, Gjnovci A, Hansotia T, Drucker DJ, Wollheim C, Burcelin R & Thorens B 2004 Gluco-incretins control insulin secretion at multiple levels as revealed in mice lacking GLP-1 and GIP receptor. *The Journal of Clinical Investigation* **113** 635-645.

Räkel A, Sheehy O, Rahme E & LeLorier J 2008 Osteoporosis among patients with type 1 and type 2 diabetes. *Diabetes & Metabolism* **34** 193-205.

Ratner RE, Dickey R, Fineman M, Maggs DG, Shen L, Strobel SA, Weyer C & Kolterman OG 2004 Amylin replacement with pramlintide as an adjunct to insulin therapy improves long-term glycaemic and weight control in Type 1 diabetes mellitus: a 1-year, randomized controlled trial. *Diabetic Medicine* **21** 1204-1212.

Reddy GK, Stehno-Bittel L, Hamade S & Enwemeka CS 2001 The biomechanical integrity of bone in experimental diabetes. *Diabetes Research and Clinical Practice* **54** 1-8.

Rees DA & Alcolado JC 2005 Animal models of diabetes mellitus. *Diabetic Medicine* **22** 359-370.

Reimer M, Holst J & Ahren B 2002 Long-term inhibition of dipeptidyl peptidase IV improves glucose tolerance and preserves islet function in mice. *European Journal of Endocrinology* **146** 717-727.

Rho J, Tsui TY & Pharr GM 1997 Elastic properties of human cortical and trabecular lamellar bone measured by nanoindentation. *Biomaterials* **18** 1325-1330.

Robey PG, Young ME, Flanders KC, Roche NS, Kondaiah P, Reddi AH, Termine JD, Sporn MB & Roberts AB 1987 Osteoblasts Synthesize and respond to transforming growth factor-type beta (TGF-beta) in vitro. *The Journal of Cell Biology* **105** 457-463.

Roschger P, Fratzl P, Eschberger J & Klaushofer K 1998 Validation of quantitative backscatterd electron imaging for the measurement of mineral density distribution in human bone biopsies. *Bone* **23** 319-326.

Rosen CJ 2004 Insulin-like growth factor I and bone mineral density: experience from animal models and human observational studies. *Best Practice & Research Clinical Endocrinology & Metabolism* **18** 423-435.

Rosen CJ & Bouxsein ML 2006 Mechanisms of disease : is osteoporosis the obesity of bone?. *Nature Clininal Practice Rheumatology* **2** 35-43.

Rosenstock J, Brazg R, Andryuk PJ, Lu K & Stein P 2006 Efficacy and safety of the dipeptidyl peptidase-4 inhibitor sitagliptin added to ongoing pioglitazone therapy in patients with type 2 diabetes: A 24-week, multicenter, randomized, double-blind, placebo-controlled, parallel-group study. *Clinical Therapeutics* **28** 1556-1568.

Rosenstock J, Foley JE, Rendell M, Landin-Olsson M, Holst JJ, Deacon CF, Rochotte E & Baron MA 2008 Effects of the dipeptidyl peptidase-IV inhibitor Vildagliptin on incretin hormones, islet function, and postprandial glycemia in subjects with impaired glucose tolerance. *Diabetes Care* **31** 30-35.

- Rosenstock J, Reusch J, Bush M, Yang F & Stewart M 2009 Potential of Albiglutide, a Long-Acting GLP-1 Receptor Agonist, in Type 2 Diabetes. *Diabetes Care* **32** 1880-1886.
- Rosenstock J, Balas B, Charbonnel B, Bolli GB, Boldrin M, Ratner R & Balena R 2013 The fate of Taspoglutide, a weekly GLP-1 receptor agonist, versus twice-daily Exenatide for type 2 diabetes. *Diabetes Care* **36** 498-504.
- Roszer T, Jozsa T, Kiss-Toth ED, Clerck ND & Balogh L 2014 Leptin receptor deficient diabetic (*db/db*) mice are compromised in postnatal bone regeneration. *Cell and Tissue Research* **356** 195-206.
- Rouille Y, Bianchi M, Irminger J & Halban PA 1997a Role of the prohormone convertase PC2 in the processing of proglucagon to glucagon. *FEBS Letters* **413** 119-123.
- Rouille Y, Kantengwa S, Irminger J & Halban PA 1997b Role of the prohormone convertase PC3 in the processing of proglucagon to glucagon-like peptide 1. *The Journal of Biological Chemistry* **272** 32810–32816.
- Rupert G, Yip C, Boylan MO, Kieffer TJ & Wolfe MM 1998 Functional GIP receptors are present on adipocytes. *Endocrinology* **139** 4004-4007.
- Ryan GJ, Jobe LJ & Martin R 2005 Pramlintide in the treatment of type 1 and type 2 diabetes mellitus. *Clinical Therapeutics* **27** 1500-1512.
- Rzonca SO, Suva LJ, Gaddy D, Montague DC & Lecka-Czernik B 2004 Bone is a target for the antidiabetic compound Rosiglitazone. *Endocrinology* **145** 401-406.
- Saito M & Marumo K 2010 Collagen cross-links as a determinant of bone quality : a possible explanation for bone fragility in aging, osteoporosis and diabetes mellitus. *Osteoporosis International* **21** 195-214.

Salera M, Giacomoni P, Pironi L, Cornia G, Capelli M, Marini A, Benfenati F, Miglioli M & Barbara L 1982 Gastric inhibitory polypeptide release after oral glucose: relationship to glucose intolerance, diabetes mellitus, and obesity. *The Journal of Clinical Endocrinology and Metabolism* **55** 329-336.

Sandhu MS, Heald AH, Gibson JM, Cruickshank JK, Dunger DB & Wareham NJ 2002 Circulating concentrations of insulin-like growth factor-I and development of glucose intolerance: a prospective observational study. *The Lancet* **359** 1740-1745.

Schjoldager BTG, Baldissera FGA, Mortensen PE, Holst JJ & Christiansen J 1988 Oxyntomodulin: a potential hormone from the distal gut. Pharmacokinetics and effects on gastric acid and insulin secretion in man. *European Journal of Clinical Investigation* **18** 499-503.

Schwartz AV, Sellmeyer DE, Ensrud KE, Cauley JA, Tabor HK, Schreiner PJ, Jamal SA, Black DM & Cummings SR 2000 Older women with diabetes have an increased risk of fracture: a prospective study. *The Journal of Clinical Endocrinology & Metabolism* **86** 32-38.

Schwartz AV, Hillier TA, Sellmeyer DE, Resnick HE, Gregg E, Ensrud KE, Schreiner PJ, Margolis KL, Cauley JA, Nevitt MC, Black DM & Cummings SR 2002 Older women with diabetes have a higher risk of falls. *Diabetes Care* **25** 1749-1754.

Schwartz AV, Sellmeyer DE, Vittinghoff E, Palermo L, Lecka-Czernik B, Feingold KR, Strotmeyer ES, Resnick HE, Carbone L, Beamer BA, Park SW, Lane NE, Harris TB & Cummings SR 2006 Thiazolidinedione use and bone loss in older diabetic adults. *The Journal of Clinical Endocrinology & Metabolism* **91** 3349-3354.

Scrocchi LA, Brown TJ, Maclusky N, Brubaker PL, Auerbach AB, Joyner AL & Drucker DJ 1996 Glucose intolerance but normal satiety in mice with a null

mutation in the glucagon-like peptide 1 receptor gene. *Nature Medicine* **2** 1254-1258.

Sebokova E, Christ AD, Wang H, Sewing S, Dong JZ, Taylor J, Cawthorne MA & Culler MD 2010 Taspoglutide, an analog of human glucagon-like peptide-1 with enhanced stability and *in vivo* potency. *Endocrinology* **151** 2474 –2482.

Seeman E 2002 Pathogenesis of bone fragility in women and men. *Lancet* **359** 1841–1850.

Seeman E & Delmas PD 2006 Bone quality - the material and structural basis of bone strength and fragility. *The New England Journal of Medicine* **354** 2250-2261.

Sondergaard BC, Wulf H, Henriksen K, Schaller S, Oestergaard S, Qvist P, Tankó LB, Bagger YZ, Christiansen C & Karsdal MA 2006 Calcitonin directly attenuates collagen type II degradation by inhibition of matrix metalloproteinase expression and activity in articular chondrocytes. *Osteoarthritis and Cartilage* **14** 759-768.

Sorocéanu MA, Miao D, Bai X, Su H, Goltzman D & Karaplis AC 2004 Rosiglitazone impacts negatively on bone by promoting osteoblast/osteocyte apoptosis. *Journal of Endocrinology* **183** 203-216.

Steppan CM, Crawford DT, Chidsey-Frink KL, Ke HZ & Swick AG 2000 Leptin is a potent stimulator of bone growth in ob/ob mice. *Regulatory Peptides* **92** 73-78.

Stonehouse AH, Darsow T & Maggs DG 2012 Incretin-based therapies. *Journal of Diabetes* **4** 55-67.

Stumvoll M, Goldstein BJ & van Haeften TW 2005 Type 2 diabetes: principles of pathogenesis and therapy. *The Lancet* **365** 1333-1346.

- Sudo H, Kodama H, Amagai Y, Yamamoto S & Kasai S 1983 In vitro differentiation and calcification in a new clonal osteogenic cell line derived from newborn mouse calvaria. *The Journal of Cell Biology* **96** 191-198.
- Tahrani AA, Bailey CJ, Del Prato S & Barnett AH 2011 Management of type 2 diabetes: new and future developments in treatment. *The Lancet* **378** 182-197.
- Takayanagi H, Kim S, Matsuo K, Suzuki H, Suzuki T, Sato K, Yokochi T, Odak H, Nakamurak K, Ida N, Wagner EF & Taniguch T 2002 RANKL maintains bone homeostasis through c-FOS-dependent induction of interferon-beta. *Nature* **416** 744-749.
- Tamasi JA, Arey BJ, Bertolini DR & Feyen JHM 2003 Characterization of Bone Structure in Leptin Receptor-Deficient Zucker (*fa/fa*) Rats. *Journal of Bone and Mineral Research* **18** 1605-1611.
- Tamura K, Minami K, Kudo M, Iemoto K, Takahashi H & Seino S 2015 Liraglutide improves pancreatic beta cell mass and function in alloxan-induced diabetic mice. *Plos One* **10** e0126003.
- Tang Y, Wu X, Lei W, Pang L, Wan C, Shi Z, Zhao L, Nagy TR, Peng X, Hu J, Feng X, Hul WV, Wan M & Cao X 2009 TGF- β 1-induced migration of bone mesenchymal stem cells couples bone resorption with formation. *Nature Medicine* **15** 757-.
- Tartaglia LA, Dembski M, Weng X, Deng N, Culpepper J, Devos R, Richards GJ, Campfield LA, Clark FT, Deeds J, Muir C, Sanker S, Moriarty A, Moore KJ, Smutko JS, Mays GG, Wool EA, Monroe CA & Tepper RI 1995 Identification and expression cloning of a leptin receptor, OB-R. *Cell* **83** 1263-1271.
- Teppala S & Shankar A 2010 Association between serum IGF-1 and diabetes among US adults. *Diabetes Care* **33** 2257-2259.

Tharakan G, Tracia T & Bloom S 2011 Emerging therapies in the treatment of 'diabetes': beyond GLP-1. *Trends in Pharmacological Sciences* **32** 8-15.

The Jackson Laboratory 2015 Genetics and genomics [Online].
<https://www.jax.org/genetics-and-healthcare/genetics-and-genomics/why-mouse-genetics#>

Thomas T, Gori F, Khosla S, Jensen MD, Burguera B & Riggs BL 1999 Leptin acts on human marrow stromal cells to enhance differentiation to osteoblasts and to inhibit differentiation to adipocytes. *Endocrinology* **140** 1630-1638.

Thomas DM, Udagawa N, Hards DK, Quinn JMW, Moseley JM, Findlay DM & Best JD 1998 Insulin receptor expression in primary and cultured osteoclast-like cells. *Bone* **23** 181-186.

Thorens B 1992 Expression cloning of the pancreatic J8 cell receptor for the glucoretin hormone glucagon-like peptide 1. *Proceedings of the National Academic of Sciences of the United States of America* **89** 8641-8645.

Thorens B, Porret A, Buhler L, Deng S, Morel P & Widmann C 1993 Cloning and functional expression of the human islet GLP-1 receptor demonstration that Exendin-4 is an agonist and Exendin-(9-39) an antagonist of the receptor. *Diabetes* **42** 1678-1682.

Thrailkill KM, Lumpkin CK, Bunn RC, Kemp SF & Fowlkes JL 2005a Is insulin an anabolic agent in bone? Dissecting the diabetic bone for clues. *American Journal of Physiology - Endocrinology and Metabolism* **289** E735-E745.

Thrailkill KM, Liu L, Wahl EC, Bunn RC, Perrien DS, Cockrell GE, Skinner RA, Hogue WR, Carver AA, Fowlkes JL, Aronson J & Lumpkin CK 2005b Bone formation is impaired in a model of type 1 diabetes. *Diabetes* **54** 2875-2881.

Trumper A, Trumper K, Trusheim H, Arnold R, Goke B & Horsch D 2001 Glucose-dependent insulintropic polypeptide is the growth factor for beta (INS-1) cells by pleiotropic signaling. *Molecular Endocrinology* **15** 1559-1570.

Trumper A, Trumper K & Horsch D 2002 Mechanisms of mitogenic and anti-apoptotic signaling by glucose-dependent insulintropic polypeptide in beta (INS-1)-cells. *Journal of Endocrinology* **174** 233-246.

Tsukiyama K, Yamada Y, Yamada C, Harada N, Kawasaki Y, Ogura M, Bessho K, Li M, Amizuka N, Sato M, Udagawa N, Takahashi N, Tanaka K, Oiso Y & Seino Y 2006 Gastric inhibitory polypeptide as an endogenous factor promoting new bone formation after food ingestion. *Molecular Endocrinology* **20** 1644-1651.

Turner CH 2006 Bone strength : current concepts. *Annals of the New York Academy of Sciences* **1068** 429-446.

Turner PV, Brabb T, Pekow C & Vasbinder MA 2011 Administration of substances to laboratory animals: routes of administration and factors to consider. *Journal of the American Association for Laboratory Animal Science* **50** 600-613.

Ugleholdt R, Poulsen MH, Holst PJ, Irminger J, Orskov C, Pedersen J, Rosenkilde MM, Zhu X, Steiner DF & Holst JJ 2006 Prohormone Convertase 1/3 Is Essential for Processing of the Glucose-dependent Insulintropic Polypeptide Precursor. *Journal of Biological Chemistry* **281** 11050-11057.

Usdin TB, Mezey E, Button DC, Brownstein MJ & Bonner TI 1993 Gastric inhibitory polypeptide receptor, a member of the secretin-vasoactive intestinal peptide receptor family, is widely distributed in peripheral organs and the brain. *Endocrinology* **133** 2861-2870.

Valeriya L, Roberto L, Piero M, Silvia DG, Marju O, Peter A, Marketa S, Charlotte L, Karl-Fredrik E, Asa-Linda L, Rita M, Goran B, Tiinamaija T, Peter N,

Stefano DP & Leif G 2007 Mechanisms by which common variants in the TCF7L2 gene increase risk of type 2 diabetes. *Journal of Clinical Investigation* **117** 2155-2163.

Van't Hof RJ 2012 Analysis of bone architecture in rodents using microcomputed tomography. *Bone Research Protocols, Methods in Molecular Biology* **816** 461-476.

Vestergaard P 2007 Discrepancies in bone mineral density and fracture risk in patients with type 1 and type 2 diabetes - a meta-analysis. *Osteoporosis International* **18** 427-444.

Vestergaard P 2011 Diabetes and bone. *Journal of Diabetes & Metabolism Special issue 1* 1-7.

Vilsbøll T, Krarup T, Deacon CF, Madsbad S & Holst JJ 2001 Reduced postprandial concentrations of intact biologically active glucagon-like peptide 1 in Type 2 Diabetic Patients. *Diabetes* **50** 609-613.

Wang RN, Bouwens L & Kloppel G 1994 Beta-cell proliferation in normal and streptozotocin-treated newborn rats: site, dynamics and capacity. *Diabetologia* **37** 1088-1096.

Wang X, Cahill CM, Pineyro MA, Zhou J, Doyle ME & Egan JM 1999 Glucagon-like peptide-1 regulates the beta cell transcription factor, PDX-1, in insulinoma cells. *Endocrinology* **140** 4904-4907.

Wang Y, Perfetti R, Greig NH, Holloway HW, DeOre KA, Montrose-Rafizadeh C, Elahi D & Egan JM 1997 Glucagon-like peptide-1 can reverse the age-related decline in glucose tolerance in rats. *The Journal of Clinical Investigation* **99** 2883-2889.

Weiner S & Wagner HD 1998 The material bone : structure-mechanical function relations. *Annual Review of Materials Science* **28** 271-298.

Wei Y & Mojsov S 1996 Distribution of GLP-1 and PACAP receptors in human tissues. *Acta Physiologica Scandinavica* **157** 355-357.

Whitehouse WJ, Dyson ED & Jackson CK 1971 The scanning electron microscope in studies of trabecular bone from a human vertebral body. *Journal of Anatomy* **108** 481-496.

Wiernsperger NF & Bailey CJ 1999 The antihyperglycaemic effect of metformin. *Drugs* **58** 31-39.

Wilding JPH, Norwood P, T'joen C, Bastien A, List JF & Fiedorek FT 2009 A study of Dapagliflozin in patients with type 2 diabetes receiving high doses of insulin plus insulin sensitizers. Applicability of a novel insulin-independent treatment. *Diabetes Care* **32** 1656–1662.

Williams GA, Callon KE, Watson M, Costa JL, Ding Y, Dickinson M, Wang Y, Naot D, Reid IR & Cornish J 2011 Skeletal phenotype of the leptin receptor-deficient *db/db* mouse. *Journal of Bone and Mineral Research* **26** 1698-1709.

Wright A, Burden ACF, Paisey RB, Cull CA & Holman RR 2002 Sulfonylurea Inadequacy: Efficacy of addition of insulin over 6 years in patients with type 2 diabetes in the U.K. Prospective Diabetes Study (UKPDS 57). *Diabetes Care* **25** 330-336.

Xian L, Wu X, Pang L, Lou M, Rosen CJ, Qiu T, Crane J, Frassica F, Zhang L, Rodriguez JP, Jia X, Yakar S, Xuan S, Efstratiadis A, Wan M & Cao X 2012 Matrix IGF-1 maintains bone mass by activation of mTOR in mesenchymal stem cells. *Nature Medicine* **18** 1095-.

Xie D, Cheng H, Hamrick M, Zhong Q, Ding K, Correa D, Williams S, Mulloy A, Bollag W, Bollag RJ, Runner RR, McPherson JC, Insogna K & Isaacs CM 2005 Glucose-dependent insulintropic polypeptide receptor knockout mice have altered bone turnover. *Bone* **37** 759-769.

- Xie D, Zhong Q, Ding K, Cheng H, Williams S, Correa D, Bollag WB, Bollag RJ, Insogna K, Troiano N, Coady C, Hamrick M & Isaacs CM 2007 Glucose-dependent insulintropic peptide-overexpressing transgenic mice have increased bone mass. *Bone* **40** 1352-1360.
- Xu G, Stoffers DA, Habener JF & Bonner-Weir S 1999 Exendin-4 stimulates both beta-cell replication and neogenesis, resulting in increased beta-cell mass and improved glucose tolerance in diabetic rats. *Diabetes* **48** 2270-2276.
- Yabe D & Seino Y 2011 Two incretin hormones GLP-1 and GIP: Comparison of their actions in insulin secretion and β cell preservation. *Progress in Biophysics and Molecular Biology* **107** 248-256.
- Yamada C, Yamada Y, Tsukiyama K, Yamada K, Udagawa N, Takahashi N, Tanaka K, Drucker DJ, Seino Y & Inagaki N 2008 The murine glucagon-like peptide-1 is essential for control of bone resorption. *Endocrinology* **149** 574-579.
- Yamagishi S 2011 Role of advanced glycation end products (AGEs) in osteoporosis in diabetes. *Current Drug Targets* **12** 2096-2102.
- Yaturu S, Bryant B & Jain SK 2007 Thiazolidinedione Treatment Decreases Bone Mineral Density in Type 2 Diabetic Men *Diabetes Care* **30** 1574-1576.
- Yip RG, Boylan MO, Kieffer TJ & Wolfe MM 1998 Functional GIP Receptors Are Present on Adipocytes. *Endocrinology* **139** 4004-4007.
- Yki-Jarvinen H 2004 Thiazolidinediones. *The New England Journal of Medicine* **351** 1106-18.
- Yoon J & Jun H 2005 Autoimmune destruction of pancreatic beta-cells. *American Journal of Therapeutics* **12** 580-591.
- Zemmit P, Alberti KGMM & Shaw J 2001 Global and societal implications of the diabetes epidemic. *Nature* **414** 782-787.

- Zhang L, Bruce-Keller AJ, Dasuri K, Nguyen AT, Liu Y & Keller JN 2009 Diet-induced metabolic disturbances as modulators of brain homeostasis. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1792** 417-422.
- Zhang L, Nakayama M & Eisenbarth GS 2008 Insulin as an autoantigen in NOD/human diabetes. *Current Opinion in Immunology* **20** 111-118.
- Zhang M, Xuan S, Bouxsein ML, von Stechow D, Akeno N, Faugere MC, Malluche H, Zhao G, Rosen CJ, Efstratiadis A & Clemens TL 2002 Osteoblast-specific knockout of the insulin-like growth factor (IGF) receptor gene reveals an essential role of IGF signaling in bone matrix mineralization. *Journal of Biological Chemistry* **277** 44005-44012.
- Zhong Q, Ding K, Mulloy AL, Bollag RJ & Isales CM 2003 Glucose-dependent insulinotropic peptide stimulates proliferation and TGF- β release from MG-63 cells. *Peptides* **24** 611-616.
- Zhong Q, Itokawa T, Sridhar S, Ding K, Xie D, Kang B, Bollag WB, Bollag RJ, Hamrick M, Insogna K & Isales CM 2007 Effects of glucose-dependent insulinotropic peptide on osteoclast function. *American Journal of Physiology - Endocrinology And Metabolism* **292** E543-E548.
- Zhukouskaya VV, Shepelkevich AP & Chiodini I 2014 Bone health in type 1 diabetes: where we are now and how we should proceed. *Advances in Endocrinology* <http://dx.doi.org/10.1155/2014/982129>.
- Zoetis T, Tassinari MS, Bagi C, Walthall K & Hurtt ME 2003 Species comparison of postnatal bone growth and development. *Birth Defects Research* **68** 86-110.